



Liu, Z., Dumville, J. C., Norman, G., Westby, M. J., Blazeby, J., McFarlane, E., Welton, N. J., O'Connor, L., Cawthorne, J., George, R. P., Crosbie, E. J., Rithalia, A. D., & Cheng, H. Y. (2018). Intraoperative interventions for preventing surgical site infection: An overview of Cochrane Reviews. *Cochrane Database of Systematic Reviews*, 2018, [CD012653.pub2].  
<https://doi.org/10.1002/14651858.CD012653.pub2>

Publisher's PDF, also known as Version of record

Link to published version (if available):  
[10.1002/14651858.CD012653.pub2](https://doi.org/10.1002/14651858.CD012653.pub2)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at <http://cochranelibrary-wiley.com/doi/10.1002/14651858.CD012653.pub2/abstract> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



**Cochrane**  
**Library**

**Cochrane** Database of Systematic Reviews

## **Intraoperative interventions for preventing surgical site infection: an overview of Cochrane Reviews (Review)**

Liu Z, Dumville JC, Norman G, Westby MJ, Blazeby J, McFarlane E, Welton NJ, O'Connor L, Cawthorne J, George RP, Crosbie EJ, Rithalia AD, Cheng HY

Liu Z, Dumville JC, Norman G, Westby MJ, Blazeby J, McFarlane E, Welton NJ, O'Connor L, Cawthorne J, George RP, Crosbie EJ, Rithalia AD, Cheng HY.

Intraoperative interventions for preventing surgical site infection: an overview of Cochrane Reviews.

*Cochrane Database of Systematic Reviews* 2018, Issue 2. Art. No.: CD012653.

DOI: 10.1002/14651858.CD012653.pub2.

**[www.cochranelibrary.com](http://www.cochranelibrary.com)**

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
BACKGROUND . . . . .	3
OBJECTIVES . . . . .	5
METHODS . . . . .	6
RESULTS . . . . .	9
Figure 1. . . . .	10
DISCUSSION . . . . .	18
AUTHORS' CONCLUSIONS . . . . .	20
ACKNOWLEDGEMENTS . . . . .	21
REFERENCES . . . . .	21
ADDITIONAL TABLES . . . . .	25
APPENDICES . . . . .	123
CONTRIBUTIONS OF AUTHORS . . . . .	128
DECLARATIONS OF INTEREST . . . . .	129
SOURCES OF SUPPORT . . . . .	129

# Intraoperative interventions for preventing surgical site infection: an overview of Cochrane Reviews

Zhenmi Liu<sup>1</sup>, Jo C Dumville<sup>1</sup>, Gill Norman<sup>1</sup>, Maggie J Westby<sup>1</sup>, Jane Blazeby<sup>2</sup>, Emma McFarlane<sup>3</sup>, Nicky J Welton<sup>2</sup>, Louise O'Connor<sup>4</sup>, Julie Cawthorne<sup>4</sup>, Ryan P George<sup>4</sup>, Emma J Crosbie<sup>5</sup>, Amber D Rithalia<sup>6</sup>, Hung-Yuan Cheng<sup>7</sup>

<sup>1</sup>Division of Nursing, Midwifery and Social Work, School of Health Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK. <sup>2</sup>NIHR Bristol Biomedical Research Centre, School of Social and Community Medicine, Bristol Medical School, University of Bristol, Bristol, UK. <sup>3</sup>Centre for Guidelines, National Institute for Health and Care Excellence, Manchester, UK. <sup>4</sup>Infection Prevention and Control / Tissue Viability Team, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK. <sup>5</sup>Division of Cancer Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK. <sup>6</sup>Independent Researcher, Leeds, UK. <sup>7</sup>Bristol Centre for Surgical Research, Bristol Medical School, University of Bristol, Bristol, UK

Contact address: Zhenmi Liu, Division of Nursing, Midwifery and Social Work, School of Health Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, Jean McFarlane Building, Oxford Road, Manchester, M13 9PL, UK. [zhenmi.liu@manchester.ac.uk](mailto:zhenmi.liu@manchester.ac.uk), [liuzhenmi1983@hotmail.com](mailto:liuzhenmi1983@hotmail.com).

**Editorial group:** Cochrane Wounds Group.

**Publication status and date:** New, published in Issue 2, 2018.

**Citation:** Liu Z, Dumville JC, Norman G, Westby MJ, Blazeby J, McFarlane E, Welton NJ, O'Connor L, Cawthorne J, George RP, Crosbie EJ, Rithalia AD, Cheng HY. Intraoperative interventions for preventing surgical site infection: an overview of Cochrane Reviews. *Cochrane Database of Systematic Reviews* 2018, Issue 2. Art. No.: CD012653. DOI: 10.1002/14651858.CD012653.pub2.

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Surgical site infection (SSI) rates vary from 1% to 5% in the month following surgery. Due to the large number of surgical procedures conducted annually, the costs of these SSIs can be considerable in financial and social terms. Many interventions are used with the aim of reducing the risk of SSI in people undergoing surgery. These interventions can be broadly delivered at three stages: preoperatively, intraoperatively and postoperatively. The intraoperative interventions are largely focused on decontamination of skin using soap and antiseptics; the use of barriers to prevent movement of micro-organisms into incisions; and optimising the patient's own bodily functions to promote best recovery. Both decontamination and barrier methods can be aimed at people undergoing surgery and operating staff. Other interventions focused on SSI prevention may be aimed at the surgical environment and include methods of theatre cleansing and approaches to managing theatre traffic.

### Objectives

To present an overview of Cochrane Reviews of the effectiveness and safety of interventions, delivered during the intraoperative period, aimed at preventing SSIs in all populations undergoing surgery in an operating theatre.

### Methods

Published Cochrane systematic reviews reporting the effectiveness of interventions delivered during the intraoperative period in terms of SSI prevention were eligible for inclusion in this overview. We also identified Cochrane protocols and title registrations for future inclusion into the overview. We searched the Cochrane Library on 01 July 2017. Two review authors independently screened search results and undertook data extraction and 'Risk of bias' and certainty assessment. We used the ROBIS (risk of bias in systematic reviews) tool to assess the quality of included reviews, and we used GRADE methods to assess the certainty of the evidence for each outcome. We summarised the characteristics of included reviews in the text and in additional tables.

## Main results

We included 32 Cochrane Reviews in this overview: we judged 30 reviews as being at low risk of bias and two at unclear risk of bias. Thirteen reviews had not been updated in the past three years. Two reviews had no relevant data to extract. We extracted data from 30 reviews with 349 included trials, totaling 73,053 participants. Interventions assessed included gloving, use of disposable face masks, patient oxygenation protocols, use of skin antiseptics for hand washing and patient skin preparation, vaginal preparation, microbial sealants, methods of surgical incision, antibiotic prophylaxis and methods of skin closure. Overall, the GRADE certainty of evidence for outcomes was low or very low. Of the 77 comparisons providing evidence for the outcome of SSI, seven provided high- or moderate-certainty evidence, 39 provided low-certainty evidence and 31 very low-certainty evidence. Of the nine comparisons that provided evidence for the outcome of mortality, five provided low-certainty evidence and four very low-certainty evidence.

There is high- or moderate-certainty evidence for the following outcomes for these intraoperative interventions. (1) Prophylactic intravenous antibiotics administered before caesarean incision reduce SSI risk compared with administration after cord clamping (10 trials, 5041 participants; risk ratio (RR) 0.59, 95% confidence interval (CI) 0.44 to 0.81; high-certainty evidence - assessed by review authors). (2) Preoperative antibiotics reduce SSI risk compared with placebo after breast cancer surgery (6 trials, 1708 participants; RR 0.74, 95% CI 0.56 to 0.98; high-certainty evidence - assessed by overview authors). (3) Antibiotic prophylaxis probably reduce SSI risk in caesarean sections compared with no antibiotics (82 relevant trials, 14,407 participants; RR 0.40, 95% CI 0.35 to 0.46; moderate-certainty evidence; downgraded once for risk of bias - assessed by review authors). (4) Antibiotic prophylaxis probably reduces SSI risk for hernia repair compared with placebo or no treatment (17 trials, 7843 participants; RR 0.67, 95% CI 0.54 to 0.84; moderate-certainty evidence; downgraded once for risk of bias - assessed by overview authors); (5) There is currently no clear difference in the risk of SSI between iodine-impregnated adhesive drapes compared with no adhesive drapes (2 trials, 1113 participants; RR 1.03, 95% CI 0.66 to 1.60; moderate-certainty evidence; downgraded once for imprecision - assessed by review authors); (6) There is currently no clear difference in SSI risk between short-term compared with long-term duration antibiotics in colorectal surgery (7 trials; 1484 participants; RR 1.05 95% CI 0.78 to 1.40; moderate-certainty evidence; downgraded once for imprecision - assessed by overview authors). There was only one comparison showing negative effects associated with the intervention: adhesive drapes increase the risk of SSI compared with no drapes (5 trials; 3082 participants; RR 1.23, 95% CI 1.02 to 1.48; high-certainty evidence - rated by review authors).

## Authors' conclusions

This overview provides the most up-to-date evidence on use of intraoperative treatments for the prevention of SSIs from all currently published Cochrane Reviews. There is evidence that some interventions are useful in reducing SSI risk for people undergoing surgery, such as antibiotic prophylaxis for caesarean section and hernia repair, and also the timing of prophylactic intravenous antibiotics administered before caesarean incision. Also, there is evidence that adhesive drapes increase SSI risk. Evidence for the many other treatment choices is largely of low or very low certainty and no quality-of-life or cost-effectiveness data were reported. Future trials should elucidate the relative effects of some treatments. These studies should focus on increasing participant numbers, using robust methodology and being of sufficient duration to adequately assess SSI. Assessment of other outcomes such as mortality might also be investigated as part of non-experimental prospective follow-up of people with SSI of different severity, so the risk of death for different subgroups can be better understood.

## PLAIN LANGUAGE SUMMARY

### Overview of Cochrane Reviews of interventions used during surgery for preventing surgical site infection

#### What is the aim of this overview of reviews?

To identify and summarise all evidence from Cochrane Reviews on interventions to prevent surgical site infections (SSIs) that are delivered while surgery is taking place (during the intraoperative period).

#### Key messages

We cannot be certain about the effectiveness in preventing SSI of the majority of intraoperative interventions, as we judged the certainty of the evidence to be generally low or very low. In some circumstances (listed below), antibiotics were effective for the prevention of SSI. There is no high- or moderate-certainty evidence for the relative effects of intraoperative interventions on mortality, and no data at all for quality of life or costs. For these reasons, we cannot be certain whether these antibiotics, which are effective at preventing SSI,

have any negative effects on mortality or quality of life. Larger trials with appropriate methods are needed to measure the outcomes that are important to both patients and health professionals.

### **What was studied in the overview?**

If bacteria get into a surgical cut during surgery, this can result in a wound infection commonly called an SSI. SSIs are one of the most common forms of healthcare-associated infections, with around 1 in 20 surgical patients developing an SSI in hospital. SSIs can also develop after people have left hospital. SSIs can result in delayed wound healing, increased hospital stays, increased use of antibiotics, unnecessary pain and, in extreme cases, death. Their prevention is therefore a key aim for health services. Many interventions are used to reduce the risk of SSI in people having surgery. These interventions can be delivered at three stages: before, during and after the operation. It is therefore important to identify interventions that can reduce the incidence of SSIs. This overview focuses only on interventions delivered during surgery.

### **What are the main results of the overview?**

In July 2017 we searched for Cochrane Reviews involving interventions for preventing SSIs during surgery. We found a total of 32 Cochrane Reviews that could be included in this overview. Two reviews had no relevant data to extract so we extracted data from 30 reviews with 349 included trials, totaling 73,053 participants. Interventions assessed included use of disposable face masks and surgical gloves, the use of oxygen during surgery, antiseptics for hand washing, patient skin preparation and cleaning the vagina before caesarean section, methods of surgical incision and skin closure and use of antibiotics to prevent infection.

Evidence of at least moderate certainty indicates that the following interventions reduce SSI risk: (1) antibiotics administered via drip before caesarean incision reduce SSI risk compared with administration after cord clamping (high-certainty evidence); (2) giving antibiotics before surgery reduces SSI risk compared with placebo after breast cancer surgery (high-certainty evidence); (3) antibiotics used to prevent wound infections probably reduce SSIs for caesarean section compared with no antibiotics (moderate-certainty evidence); (4) antibiotics used to prevent wound infections probably reduce SSI risk for hernia repair compared with placebo or no treatment (moderate-certainty evidence); (5) iodine-impregnated adhesive drapes probably make no difference to SSI risk compared with no adhesive drapes (moderate-certainty evidence); (6) there is probably no difference in SSI risk when antibiotics are given in the short-term compared to the long-term during colorectal surgery (moderate-certainty evidence). One comparison showed that adhesive drapes increase the SSI risk compared with no drapes (high-certainty evidence). Overall, we judged the certainty of evidence for our primary outcomes (SSIs and death) to be low or very low.

Clinicians can use the evidence summarised in this overview to choose the best intervention for people having surgery. However, many of the comparisons were supported by low- or very low-certainty evidence and so require further evidence to support future decision making. This overview can also be used by policymakers in developing local and regional protocols or guidelines and can reveal knowledge gaps for future research.

### **How up to date is this overview?**

We searched for reviews that had been published up to July 2017. Of the 32 reviews included in this overview, 13 reviews had not been updated in the past three years.

## **BACKGROUND**

### **Description of the condition**

Millions of surgical procedures are conducted around the world each year. Most procedures result in surgical wounds that heal by primary intention, where wound edges are re-approximated using

sutures, staples, clips or glue. Some surgical wounds are left open to heal (where closure is not appropriate because of infection, physical impossibility of approximating wound edges or because of the need to allow drainage) and some wounds break down following closure; these open wounds heal from the 'bottom-up' (known as 'healing by secondary intention').

Surgical wounds are at risk from microbial contamination and thus possible infection. Contamination may originate from the

patient, for example when microbes on the skin enter a wound, or from the surrounding environment, for example from operating staff, the theatre, or wider hospital and home environments. SSIs are relatively common: a recent US study with assessment in 183 hospitals involving 11,282 patients found that 452 people (4%) developed hospital-acquired infection; of these, 21.8% were SSIs (Magill 2014). Similar SSI incidence estimates have been reported in France (Astagneau 2009). In the UK around 2% to 5% of surgical patients develop SSIs (NICE 2008; Public Health England 2014) although the percentage varies greatly depending on the circumstances, including the contamination level of the surgery. In England, a 2006 survey of hospital-acquired infections reported that 8% of patients in hospitals had an infection while an inpatient, of which 14% were considered SSIs (Hospital Infection Society 2007; Smyth 2008). Many quoted incidence estimates for SSI are likely to be underestimates because infections that developed outside hospitals were not considered (Bruce 2001; Gibbons 2011). While more data are available for Western healthcare settings, SSI was identified as the leading cause of hospital-acquired infection in a systematic review of studies in low- and middle-income countries (Allegranzi 2010).

SSI is a serious global issue that can lead to significant morbidity, need for re-intervention and treatment (including antibiotic use), delayed wound healing, and in very serious infections, the possibility of death (Awad 2012; Brown 2014; CDC 2017). SSIs also increase consumption of healthcare resources. Recent figures from the UK suggest that SSIs lead to a median increased hospital stay of 10 days (95% confidence interval (CI) 7 to 13 days) with an associated median additional cost attributed to SSI of GBP 5239 (95% CI GBP 4622 to 6719) (Jenks 2014). The UK National Institute for Health and Care Excellence (NICE) identified that an SSI increased the costs of surgery by two to five times (NICE 2008). In the USA, De Lissovoy 2009 estimated that the extended length of stay and increased treatment costs associated with SSIs over a one-year period led to approximately 1 million additional inpatient-days, costing an additional USD 1.6 billion.

### SSI risk

A patient's overall physical health can predict the risk of SSI, as can the type of surgical procedure (in terms of potential for contamination) and duration of surgery. These factors are collectively included in the National Nosocomial Infections Surveillance risk index (Gaynes 2001; SWI Task Force 1992), which proposes three criteria to assess risk: American Society of Anesthesiologists (ASA) score of 3, 4, or 5 (ASA 2014); wound class (see below); and duration of surgery. Other risk factors for SSI are suggested; such as if surgery is elective or emergency, but supporting data for these risk factors are more limited.

### Wound class

Wound class is assessed using the classification system adopted by the Centres for Disease Control and Prevention (HICPAC 1999).

- Clean: non-infective operative wounds in which no inflammation is encountered, and neither the respiratory, alimentary, genito-urinary tract nor the oropharyngeal cavity is entered. In addition these cases are elective, have primary closure, and wounds are drained with closed drainage systems when required.
- Clean/contaminated: operative wounds in which the respiratory, alimentary, genital or urinary tract is entered under controlled conditions and without unusual contamination. Specifically, operations involving the biliary tract, appendix, vagina and oropharynx are included in this category, provided no evidence of infection or a major break in sterile technique is encountered.
- Contaminated: fresh, accidental wounds, operations with major breaks in sterile technique or gross spillage from the gastro-intestinal tract, and incisions in which acute, non-purulent inflammation is encountered.
- Dirty: old traumatic wounds with retained devitalised (dead) tissue and those that involve existing clinical infection or perforated viscera (internal organs or gut). This definition suggests that organisms causing postoperative infection were present in the operative field before the operation.

In the UK data from 232 NHS hospitals on 620,535 surgical procedures reported SSI rates of: 0.5% for knee prosthesis; 1% for cardiac surgery (non-coronary artery bypass graft); 0.6% for hip prosthesis and 5% for limb amputation (all clean surgery) (Health Protection Agency 2015). This is in contrast to the incidence of SSI following surgery on the large bowel (contaminated surgery) of 9.7% (Health Protection Agency 2015). Europe-wide surveillance also reports higher incidence of SSI in colon surgery (9.5% of surgeries resulting in SSI) (ECDC 2013).

### Definition of SSI

Although there is no single agreed diagnostic tool or protocol to confirm the presence of an SSI, (Bruce 2001 identified 41 different definitions for SSI and 13 grading scales), the Centers for Disease Control and Prevention (CDC) definition is commonly used (Horan 1992).

A superficial SSI is defined as: "an infection occurring within 30 days after the operation and only involving the skin and subcutaneous tissue of the incision that is associated with at least one of the following:

- purulent drainage, with or without laboratory confirmation, from the surgical site;
- organisms isolated from an aseptically-obtained culture of fluid or tissue from the surgical site;
- at least one of the following signs or symptoms of infection: pain or tenderness, localised swelling, redness or heat, and superficial incision is deliberately opened by the surgeon and is

culture-positive or not cultured. A culture-negative finding does not meet this criterion;

- diagnosis of SSI by the surgeon or attending physician.”

A deep incisional SSI is defined as: “infection that occurs within 30 days after the operative procedure if no implant is left in place, or within one year if an implant is left in place, and the infection appears to be related to the operative procedure *and* involves deep soft tissues (e.g. fibrous connective tissues and muscle layers) of the incision associated with one of the following:

- purulent drainage from the deep incision, but not from the organ/space component of the surgical site;
- a deep incision spontaneously dehisces (opens up) or is deliberately opened by the surgeon and is culture-positive or not cultured when the patient has at least one of the following symptoms: fever or localised pain or tenderness;
- an abscess, or other evidence of infection involving the deep incision is found on direct examination, during re-operation, or by histopathologic or radiologic examination;
- diagnosis of a deep incisional SSI by a surgeon or attending physician.”

## Description of the interventions

Many interventions are used with the aim of reducing the risk of SSI in people undergoing surgery. These interventions can be delivered at three stages: preoperatively, intraoperatively and post-operatively (Goodman 2017). For the purpose of this review we define:

- the preoperative phase as the time period between the decision for the need for surgery and when everything is ready for the operation to start, that is, the patient is on the operating table (for this review we have assumed that staff are ready to proceed with surgery at this point - thus the preparation of operative staff occurs in this preoperative period);
- the intraoperative phase is the time period from when the patient is on the operating table to when the operation has finished and the wound is closed (if relevant). We consider any activity taking place after induction of anaesthesia to be in this phase because this starts in the operating theatre itself. For this review, where it is clear that antibiotics were given very soon before the incision, we consider this to be intraoperative, that is, prophylactic intravenous antibiotics administered less than 60 minutes before surgery;
- the postoperative phase as the time period from the end of the intraoperative phase to resolution of surgical procedure (which we acknowledge could take several, weeks or months for some patients). We note that whilst dressings, wound drains and negative pressure wound therapy are often placed over wounds at the end of surgery, their use is predominantly outside of theatre, so they are considered in the postoperative phase.

Table 1 details key intervention types used at each stage of the operative pathway, but is not an exhaustive list. Most interventions listed are probably independent of each other and would generally be delivered concurrently. However, the interventions listed could also be grouped together as a care bundle, where a care bundle is defined as a group of three to five evidence-based interventions that are delivered together.

This overview of reviews will focus on interventions delivered in the intraoperative phase.

## How the intervention might work

See Table 1. The interventions are largely focused on decontamination of skin using soap and antiseptics; the use of barriers to prevent movement of micro-organisms into wounds; and optimising the patient's own bodily functions to promote best recovery. Both decontamination and barrier methods can be aimed at people undergoing surgery and operating staff. Other interventions focused on SSI prevention may be aimed at the surgical environment and include methods of theatre cleansing and approaches to theatre traffic (i.e. how the movement of staff in and out of theatre is managed).

## Why it is important to do this overview

The *Cochrane Handbook for Systematic Reviews of Interventions* describes a Cochrane overview of reviews as being “intended primarily to summarize multiple Cochrane Intervention reviews addressing the effects of two or more potential interventions for a single condition or health problem” (Becker 2011).

SSIs are a prevalent problem for global healthcare and their prevention is a major focus for healthcare providers internationally. There are several Cochrane Reviews that draw together randomised controlled trial evidence for individual interventions for the prophylaxis of SSIs along the preoperative, intraoperative and postoperative pathway. Findings from these reviews have not been collated, so a transparent and usable synthesis of this evidence is required. This overview will aid decision makers aiming to draw together Cochrane evidence that spans the SSI prevention pathway. It will also be a useful resource for guideline developers, especially for the key NICE guidelines, which have not been fully updated for several years (NICE 2008). (A planned update of the guidelines was announced in 2017.) This overview will also complement other guidelines such as those produced by the World Health Organization (Allegranzi 2016a; Allegranzi 2016b).

## OBJECTIVES

To present an overview of Cochrane Reviews of the effectiveness and safety of interventions delivered during the intraoperative



period aimed at preventing SSIs in all populations undergoing surgery in an operating theatre.

## METHODS

### Criteria for considering reviews for inclusion

#### Types of studies

We included reviews published in the *Cochrane Database of Systematic Reviews* that examine the effectiveness of interventions aimed at preventing SSIs. We did not consider non-Cochrane reviews. We only included systematic reviews of randomised controlled trial (RCT) evidence for patient-focused interventions. If reviews included other study designs alongside RCTs (e.g. controlled clinical trials, quasi-randomised controlled trials, or both) we only investigated if RCT evidence was presented separately for relevant analyses (e.g. as sensitivity analyses). If so, these RCT data were included. If there were no separate data for RCTs in a review of patient-focused interventions we did not include that review in analyses. Primary RCTs published since the included reviews, but not yet included in reviews, were excluded in line with Cochrane guidance.

Where studies evaluated service-level interventions e.g. protective staff coverings, theatre traffic and environmental cleansing, designs such as interrupted time series and controlled before and after studies were more feasible and we also extracted data from these study designs as well as from RCTs (including cluster RCTs).

#### Types of participants

We included reviews of studies involving adults or children or both. We excluded reviews where inclusion criteria specified that study participants had infected wounds at baseline (i.e. treatment rather than prevention reviews). Reviews that considered both treatment and prevention studies were examined in detail to isolate relevant comparisons.

We included reviews of participants undergoing surgery of any contamination level (clean, clean/contaminated, contaminated and dirty). Reviews focused solely on graft sites and wounds of the mouth and eye were excluded. We included reviews looking at surgical wounds planned to heal by primary intention (closed wounds) and secondary intention (open wounds). Given their specialist nature, we excluded eye and oral surgeries and studies looking at infection prevention in pin sites.

#### Types of interventions

We included reviews that assessed the following interventions aimed at preventing SSIs during the intraoperative period of the patient care pathway (regardless of comparator - all were eligible):

- decontamination of patients' skin at site of surgery incision;
- use of intraoperative prophylactic antibiotics;
- skin sealants;
- use of standard and incise drapes;
- use of masks, hair covers, overshoes, gowns and other protective coverings for theatre staff;
- different glove protocols;
- use of electrosurgery for surgical incisions;
- maintaining patient homoeostasis (warming);
- maintaining patient homoeostasis (oxygenation);
- maintaining patient homoeostasis (blood glucose control);
- wound irrigation and intracavity lavage (including use of intraoperative topical antiseptics before wound closure);
- closure methods;
- theatre traffic (protocols for managing the movement of people in theatre).

We excluded reviews focusing on comparisons of different surgical approaches for the same surgery (e.g. different techniques for inguinal surgical repair; open versus closure of perianal wounds) or other interventions specific to certain types of surgery or procedures. We also excluded studies comparing different anaesthesia regimens and those investigating the use of implants or internal devices.

Where interventions were delivered at multiple time periods in the same studies, such as for assessment of antibiotics where treatment was started in one phase and continued through multiple phases (e.g. antibiotics started preoperatively and continued postoperatively), data are presented in the overview that correspond with the start of the treatment. Thus this intraoperative overview includes reviews where the start of treatment was in the intraoperative phase. Where a review contained trials that variously delivered interventions at different starting phases, we aimed to extract and present data only for those trials relevant to the intraoperative phase (that is where the treatment started in the intraoperative phase).

#### Types of outcomes

We present data according to the time points used in reviews (if reported). Where possible, we grouped data into follow-up of 30 days or less and follow-up of more than 30 days. If a review presented data at many different time points, the overview authors reported data from the time points closest to 30 days and one year, noting where other time point data were available in the original review.

#### Primary outcome

## SSIs

Occurrence of postoperative SSI as defined by the CDC criteria (Horan 1992), or the study authors' definition of SSI. Where available we present data that differentiated between superficial and deep-incisional infection.

## Secondary outcomes

### Mortality

All cause-postoperative mortality (e.g. we did not differentiate between infection-related mortality and other mortality from other causes).

### Health-related quality of life

We included quality-of-life assessments where they were reported using a validated scale that presents a single global score (e.g. SF-12, SF-36 or EQ-5D) or a validated, disease-specific questionnaire. Ideally, reported data were adjusted for baseline scores. We did not include ad hoc measures of quality of life that were not likely to be validated and would not be common to more than one trial. We did not plan to report multiple domain scores from the same measure but rather to report only overall scores for instruments e.g. physical component summary score and mental component summary score for the SF-36.

### Cost-effectiveness

Findings that considered relative costs and benefits simultaneously.

## Search methods for identification of reviews

We searched the *Cochrane Database of Systematic Reviews* (CDSR) using the search strategy presented in [Appendix 2](#). Given the large number of interventions relating to the review, the search terms focused on identification of reviews linked to SSI rather than to specific interventions. The search was undertaken on 01 July 2017 (CDSR 2017, Issue 7), after which we tracked any included reviews for updates, and followed protocols in case of full review publication until 25 July 2017 (CDSR 2017, Issue 7).

## Data collection and analysis

### Selection of reviews

Two overview authors independently screened review titles and abstracts to identify potentially relevant inclusions. We obtained

the full text of all reviews thought to be potentially eligible for further investigation. The same two overview authors independently screened the full text of all potentially relevant resources for inclusion in the overview. We recorded reasons for exclusion of any reviews excluded at this stage. Any disagreements were resolved through discussion with a third overview author. Where overview authors were also authors of included reviews we sought to avoid bias by ensuring that decisions were made by two other overview authors.

## Data extraction and management

We extracted data into a predefined and piloted data extraction form to ensure consistent data capture from each resource. Data were extracted by one overview author and independently checked by a second, with a third acting as arbiter where required. We extracted the following data for each included resource:

- study identification, review authors' details;
- review objectives;
- review inclusion and exclusion criteria;
- included settings;
- included populations, including types of surgery or procedure and depth of incision;
- all relevant comparisons and associated time points;
- concurrent intervention types that were the same for all intervention arms;
- numbers of relevant included RCTs;
- outcomes reported and details of reported outcome values;
- method and results of risk of bias and evidence quality assessment;
- GRADE assessments;
- details of any subgroup and sensitivity analyses.

Where a comparison was included in more than one review, we recorded the details multiple times (because it was relevant to each review in which it was contained). However, we reported the comparison only once for the review with the lowest risk of bias, or the most recent review if there was no difference in risk of bias assessment. We extracted meta-analysed data where possible and single study data when pooled data were not available: we extracted effect sizes with 95% confidence intervals where possible. We also extracted contextual information to enable narrative descriptions of how data were pooled (or not) presented per comparison (e.g. if some trials had been pooled for a comparison and some had not). If any information from a review was unclear or missing, we accessed the published reports of the individual trials. We did not contact study authors for details of missing data, but rather assumed that review authors had done all they could to retrieve data. We entered data into Review Manager 5 software ([RevMan 2014](#)).

## Assessment of methodological quality of included reviews

### Quality of included Cochrane Reviews

We used the risk of bias in systematic reviews (ROBIS) tool (Whiting 2016) to assess the risk of bias in systematic reviews. ROBIS assesses reviews in three phases: first, assessing relevance (optional); second, identifying concerns with the review process; and third, forming an overall judgement of the risk of bias. In the second phase, concerns with the review process fall into four domains: (1) study eligibility criteria; (2) identification and selection of studies; (3) data collection and study appraisal; and (4) synthesis and findings. Each domain contains a list of signalling questions to guide the bias assessment process. The signalling questions can be answered yes, probably yes, probably no, no or no information. Questions are worded so that a yes response relates to low concerns about the review e.g. "Did the review adhere to pre-defined objectives and eligibility criteria? and were the eligibility criteria appropriate for the review question?" At the end of each domain the assessor draws together their appraisal to indicate their concerns regarding: specification of study eligibility (domain 1); methods used to identify and select studies, or both (domain 2); methods used to collect data and appraise studies (domain 3); and the synthesis and findings (domain 4). Concerns can be graded low, high or unclear. We recorded the rationale or reasoning for decisions at each stage, that is for the signalling questions and the level of concern rated, in a table for each domain. As this overview only included Cochrane Reviews and relevance was considered as part of our screening and selection process, we did not assess relevance using the ROBIS tool (an optional first phase). Two reviewers (ZL and JD) assessed each review independently, without blinding, using a previously piloted standardised form based on the ROBIS Guidance Document and consulted each other to resolve any discordance and to compile a consensus judgment for each domain. We presented a summary of ROBIS results for each review using table format, which lent itself to presentation of data for a large number of reviews.

### Quality or certainty of evidence extracted from included reviews

It is important to present the quality or certainty of evidence from each review. We present a GRADE assessment for each eligible outcome and comparison. Where GRADE assessment was conducted in the review we extracted this assessment; however, where GRADE assessments were not available, the overview authors undertook assessment (making it clear that they had conducted the GRADE assessment post hoc).

When making decisions for the risk of bias domain, we downgraded one level when studies had been classified at high risk of bias for one or more domains and where they were classified at

unclear risk of bias for both domains that contributed to selection bias, or both.

In assessing the precision of effect estimates for SSI we followed GRADE guidance (GRADE 2013; Schünemann 2011a; Schünemann 2011b). We planned to take a conservative approach and calculated an optimal information size (OIS) for the SSI outcome using conventional sample size calculation methods and assuming a relative risk reduction of between 20% and 30% (Guyatt 2011). The OIS is summarised below but should not be treated as an optimal sample size for any future research. In GRADE assessments, the OIS is used to assess the stability of confidence intervals (CI) rather than to assess the appropriateness of a sample size to detect a difference *per se*.

Reduction in SSI from 14% to 10% (80% power; alpha 5%) = 2070 participants overall. Although on average, SSI rates are lower than 14% in many high-income countries, they can be higher in some countries and figures vary by SSI risk of the patient. We took 14% as a conservative upper estimate of SSI incidence and calculated 40% relative risk reduction.

We used the GRADE default minimum overall sample size for dichotomous outcomes of 300 in lieu of the OIS to assess precision for mortality.

If the OIS was not met we downgraded one level. We downgraded two levels if there were very few events (or very few participants for continuous outcomes). If the OIS was met we downgraded one level if the 95% CI failed to exclude important benefits and harms, which we considered as a relative risk reduction or increase of 25%.

Judgement of GRADE certainty was agreed through discussion involving at least two overview authors and involving additional overview authors where there were disagreements.

### Data synthesis

The aim of this review is to present a detailed summary of treatment-effect data for interventions aimed at SSI prevention. We present all relevant comparisons grouped by intervention type (including details of co-interventions when recorded). We also considered data according to the contamination level of surgery where possible. We use tabular formats to present summaries of treatment effects with a corresponding GRADE assessment for each comparison. Where possible we extracted meta-analysed data, along with details of model type and measures of statistical heterogeneity. Where data had not been meta-analysed we report study-level treatment effects. Results from review subgroup and sensitivity analyses are also presented. We present all data in tabular, meta-analysis or narrative formats.

Where applicable, we converted available data to risk ratios (RR). Where this was not possible we present original data. We had planned not to undertake re-analysis of data beyond conversions to RR. However, due to the inclusion of multi-stage reviews (reviews that evaluated interventions at different points on the care

pathways, i.e. pre-, intra- or postoperative, or a combination of these) we extracted data on only trials where the intervention was started in the intraoperative phase. Where it seemed appropriate for each comparison, the overview authors meta-analysed these subsets of trials. We have cautiously pooled these data into a new meta-analysis relevant to this overview of reviews, reporting the results as RR with 95% CI (again, if the subsets of trials were reported as odds ratios (OR), we converted available data to RR). We did not plan to undertake a network meta-analysis within given intervention types.


## RESULTS

See Characteristics of included reviews [Table 2](#); Characteristics of excluded reviews [Table 3](#).

### Description of included reviews

The search generated 414 records 330 of which we excluded based on the title and abstract, and 84 of which we assessed as full text. Of these, 32 reviews were eligible for this review (See [Table 2](#); [Figure 1](#)). Of the included reviews:

**Figure 1. Study flow diagram**



- two focused on theatre staff attire (Tanner 2006; Vincent 2016);
- five focused on the preparation of the surgical site (Dumville 2015; Hadiati 2014; Haas 2014; Webster 2015; Wood 2016).
- two focused on the method of surgical incision (Charoenkwan 2017; Cook 2014).
- five focused on patient homeostasis during surgery (Buchleitner 2012; Campbell 2015; Grocott 2012; Kao 2009; Wetterslev 2015). Of these, two were multi-stage reviews (that is they included trials evaluating intraoperative interventions as well as pre- or postoperative interventions, or both). In these reviews we extracted the relevant trials focusing on intraoperative intervention delivery for this review) (Buchleitner 2012; Grocott 2012).
- 12 reviews focused on the use of intraoperative prophylactic antibiotics for preventing SSIs (Gurusamy 2011; Gurusamy 2013; Gyte 2014; Jones 2014; Lipp 2013; Low 2012; Mackeen 2014; Nabhan 2016; Nelson 2014; Sanabria 2010; Sanchez-Manuel 2012; Smaill 2014). Most of these were multi-stage reviews and again, we extracted only data from trials delivering interventions that started in the intraoperative phase.
- six reviews focused on interventions for wound closure (AL-Khamis 2010; Biancari 2010; Dumville 2014; Gurusamy 2014a; Gurusamy 2014b; Mackeen 2012).

SSI was reported in 75% (24/32) of the included reviews; mortality was reported in 19% (6/32) and health-related quality of life or cost-effectiveness were not reported in any included review. Six per cent (2/32) of the reviews, reported no outcome data relevant to this overview (Campbell 2015; Low 2012). In total we extracted data from 30 reviews with 349 included trials, totaling 73,053 participants. We present SSI outcome data for 77 comparisons and mortality data for nine comparisons. Of the 52 excluded full-text reviews, 48 focused on interventions only relating to the pre- or postoperative phase (or both) and four titles were at the protocol stage only (see Table 3; Figure 1).

## Methodological quality of included reviews

### ROBIS quality of included reviews

We rated the quality of included reviews using the ROBIS tool signalling questions (Table 4; detailed assessments by signalling questions are shown in Appendix 3) presenting the overall 'Risk of bias' assessment results for each review in Table 4. Our judgements of the four domain assessment findings were as follows:

- we judged study eligibility to be at low concern for all included reviews;

- we judged the process of study identification and selection to be at low concern for all included reviews;
- for data collection and study appraisal, we judged 91% of included reviews to be at low concern. We deemed study quality not fully assessed in four studies (Cook 2014; Low 2012; Tanner 2006; Vincent 2016) and we judged these as unclear;
- for synthesis and findings, we judged 97% (31/32) of included reviews to be at low concern due to the synthesis being unlikely to produce biased results. Only one review (Cook 2014) did not consider clinical diversity across studies and bias was not explicitly addressed in the synthesis; we judged this review at high concern.

### Overall risk of bias

We considered issues around risk of bias in all reviews. In terms of the overall 'Risk of bias' assessment, we judged 94% (30/32) of reviews to be at low risk of bias.

### Quality of evidence in included reviews

Of the 32 included reviews, 31% (10/32) reported a GRADE assessment for the SSI outcome, whilst only one review (3%) reported a GRADE assessment for the mortality outcome. No other GRADE assessments were reported in the included reviews. The overview authors undertook GRADE assessment of relative treatment-effect data where no review-level GRADE assessment was available. Overall, the GRADE certainty of evidence was low or very low, as summarised in Table 5. Of the 77 comparisons presenting SSI data, we judged 51% (39/77) as low certainty and 40% (31/77) as very low certainty. Of the nine comparisons presenting mortality data, we judged 56% (5/9) as being at low certainty and 44% (4/9) as very low. Common reasons for downgrading the certainty of evidence were risk of bias of included studies, imprecision and inconsistency.

## Effect of interventions

### Analysis of results

A detailed presentation of relative treatment-effect data and GRADE assessments for all individual comparisons are in Table 5 and Table 6. Below we present a narrative summary of key findings in an order of the process of the surgery. Where included reviews contained trials investigating only intraoperative phase interventions, we interpreted results using data reported in the review, and did not return to the original studies. Where data were reported as RR, with or without 95% CIs, we

used the results directly from the reviews. Where data were reported as OR, we converted data to RR if appropriate. When the evidence was of very low certainty, we did not report RR in the main text but we did clarify RR (or original OR) in Table 5 and Table 6.

When reviews were multi-stage, that is they contained studies that variously started interventions at the pre-, intra- or postoperative stage (e.g. Antimicrobial prophylaxis for colorectal surgery (Nelson 2014)) we extracted data only for the trials relevant to this overview and reported cautious re-analysis of these extracted trials (see Data synthesis). Such re-analyses have been clearly marked in Table 5 and Table 6.

We present results by review, following an order relevant to the clinical pathway. We have made it clear that where general details of surgery were available we have reported these. Where there is a lack of detail this reflects the lack of information in the initial review. Further, we did not group the outcomes into follow-up of 30 days or less and follow-up of more than 30 days, as these time points were not recorded by the original reviews when the review authors did the meta-analysis.

## 1. Theatre staff attire

Two reviews investigated theatre staff attire interventions:

### 1.1. Double gloving for preventing SSIs

Tanner 2006 included two trials (125 participants) that compared double latex gloving with double latex gloving with a liner in a single comparison, however neither trial reported any SSI events (low-certainty evidence; downgraded twice for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 1.2. Disposable face masks for preventing SSIs

Vincent 2016 included three trials (2106 participants) that compared disposable face mask use with no mask in a single comparison. Due to clinical heterogeneity, the review did not pool data.

#### SSI

Available trial evidence reports no clear difference in SSI risk following use of disposable face masks compared with no mask (low-certainty evidence; downgraded for once for imprecision and once for inconsistency - assessed by overview authors).

No other outcome data relevant to the overview were reported.

## 2. Preparation of the surgical site

Six reviews reported interventions used to prepare the surgical site.

### 2.1. Skin antiseptics for preventing SSIs after clean surgery

Dumville 2015 included 13 trials (2623 participants in total) and evaluated a large number of different interventions, resulting in 12 comparisons of different types of skin antiseptic solutions and scrubs on SSI risk.

#### SSI

Available evidence largely reports no clear difference between different types of skin antiseptics on SSI risk. The certainty of evidence for the majority of these comparisons was low or very low. Data from one trial (542 participants) suggested that 0.5% chlorhexidine in methylated spirit may reduce SSI risk compared with povidone iodine paint (RR 0.47, 95% CI 0.27 to 0.82; low-certainty evidence; downgraded once for risk of bias and once for imprecision - assessed by overview authors). The review also grouped interventions together in an analysis based on whether treatments were aqueous or alcoholic. Data from six trials (1400 participants) showed no clear difference between aqueous solutions and alcoholic solutions (RR 0.77, 95% CI 0.51 to 1.17; low-certainty evidence; downgraded once for risk of bias and once for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 2.2. Skin preparation following caesarean section for preventing SSIs

Hadiati 2014 included five trials (1466 participants in total) and presented four different comparisons: one comparing drapes with no drapes (two trials with 1294 participants) and three (172 participants) comparing different skin antiseptics.

#### SSI

Available trial evidence reports no clear difference between compared treatments on SSIs risk (low- and very low-certainty evidence; variously downgraded once for risk of bias and once or twice for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 2.3. Vaginal preparation with antiseptic solutions for preventing SSIs

Haas 2014 included six trials (2205 participants) that compared antiseptic solutions with placebos in a single comparison.

#### SSI

Available trial evidence reports no clear difference between compared treatments on SSI risk (low-certainty evidence; downgraded

once for risk of bias and once for imprecision - assessed by review authors).  
No other outcome data relevant to the overview were reported.

#### 2.4. Plastic adhesive drapes for preventing SSIs

Webster 2015 included seven trials (4195 participants in total) and presented two comparisons.

##### SSI

The first comparison compared adhesive drapes with no drapes (five trials, 3082 participants) and found that use of adhesive drapes was associated with an increase in SSI risk (RR 1.23, 95% CI 1.02 to 1.48; high-certainty evidence - assessed by review authors). The second comparison compared iodine-impregnated adhesive drapes with no adhesive drapes (two trials; 1113 participants). Available trial evidence reports no clear difference in SSI risk (RR 1.03, 95% CI 0.66 to 1.60; moderate-certainty evidence; downgraded once for imprecision - assessed by review authors).  
No other outcome data relevant to the overview were reported.

#### 2.5. Microbial sealants for preventing SSIs

Wood 2016 included seven trials (859 participants in total) that compared application of cyanoacrylate microbial sealants with no microbial sealant in a single comparison.

##### SSI

Available trial evidence shows no clear difference in SSI risk between treatments (RR 0.53, 95% CI 0.24 to 1.18; low-certainty evidence; downgraded once for risk of bias and once for imprecision - assessed by overview authors).  
No other outcome data relevant to the overview were reported.

### 3. Making the surgical incision

#### 3.1. Scalpel versus electrosurgery for major abdominal incisions

Charoenkwan 2017 included 11 trials (2178 participants) comparing scalpel with electrosurgery in a single comparison.

##### SSI

Available trial evidence reports no clear difference between compared treatments on SSI risk (RR 1.07, 95% CI 0.74 to 1.54; low-

certainty evidence; downgraded for risk of bias and imprecision - assessed by review authors).  
No other outcome data relevant to the overview were reported.

#### 3.2. Scalpel versus no-scalpel incision for vasectomy

Cook 2014 included two trials (1182 participants in total) that compared scalpel versus no-scalpel incision for vasectomy. As these two trials differed in their duration of follow-up and the level of operator experience with the no-scalpel technique, the review did not pool data.

##### SSI

It is uncertain whether no-scalpel incision reduces SSI risk (very low-certainty evidence; downgraded once for risk of bias, once for imprecision and once for heterogeneity - assessed by overview authors). Based on ROBIS, we assessed this review as being at unclear risk of bias because it used a limited 'Risk of bias' assessment process; and also due to the lack of information about data synthesis. The review authors did not state why synthesis was done for only some of their outcomes.  
No other outcome data relevant to the overview were reported.

### 4. Treatment of the patient during surgery

#### 4.1. Warming of intravenous and irrigation fluids

Campbell 2015 included 24 studies (1250 participants in total).  
No outcome data relevant to the overview were reported.

#### 4.2. Intensive glycaemic control for preventing SSIs

Buchleitner 2012 included 12 trials (1403 participants in total). We categorised only two included trials (105 participants) as delivering interventions that started in the intraoperative phase.

##### SSI

We considered the reported outcome, 'infectious complications' to be synonymous with SSIs and pooled the data from these two trials (105 participants). It is uncertain whether intensive glycaemic control reduces SSI risk when compared with conventional glycaemic control (RR 0.71, 95% CI 0.22 to 2.26; very low-certainty evidence; downgraded twice for imprecision and once for inconsistency - assessed by overview authors).



### **Mortality**

We pooled the data from two trials (105 participants). It is uncertain whether intensive glycaemic control reduces mortality risk compared with conventional glycaemic control (RR 1.23, 95% CI 0.18 to 8.43; very low-certainty evidence; downgraded twice for imprecision, once for inconsistency - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### **4.3. Perioperative glycaemic control regimens for preventing SSIs**

[Kao 2009](#) included five trials (743 participants in total). We categorised only three included trials (589 participants) as delivering interventions that started in the intraoperative phase.

### **SSI**

Evidence from one trial (78 participants) showed no clear difference in SSI risk when applying intra- and postoperative strict glycaemic control (using intravenous insulin) compared with conventional glycaemic control (RR 0.48, 95% CI 0.04 to 5.03; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors). One trial (371 participants) showed no clear difference when applying strict intraoperative glycaemic control (using insulin infusion) with conventional glycaemic control (RR 0.86, 95% CI 0.30 to 2.52; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors). Another trial (140 participants) reported outcomes for pneumonia and wound infections and we did not consider these data. Due to variation in SSI outcomes we did not pool these trials in this overview.

### **Mortality**

Evidence from 1 trial (78 participants) showed no clear difference in overall mortality risk when applying intra- and postoperative strict compared with conventional glycaemic control (RR 0.81, 95% CI 0.30 to 2.20; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors). Similarly, another trial (371 participants) showed no clear difference when applying intraoperative strict compared with conventional glycaemic control (RR 9.05, 95% CI 0.49 to 166.88; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors). The potential for harm represented in the imprecision reported is important to acknowledge here.

No other outcome data relevant to the overview were reported.

### **4.4. Increased global blood flow for preventing SSIs**

[Grocott 2012](#) included 31 trials (5292 participants in total). We categorised 15 included trials (1202 participants) as delivering interventions that started in the intraoperative phase.

### **SSI**

Five included trials (353 participants) reported SSI data, which we pooled. Increased global blood flow (e.g. fluids and/or inotrope; oesophageal doppler) may reduce SSI risk compared with no treatment (RR 0.40, 95% CI 0.19 to 0.82; low-certainty evidence; downgraded once for imprecision and once for inconsistency - assessed by overview authors).

### **Mortality**

Fifteen relevant trials (1202 participants) reported mortality and we pooled these data. There was no clear difference in mortality risk following interventions to increase global blood flow compared with no treatment (RR 0.67, 95% CI 0.40 to 1.13; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### **4.5. High perioperative inspiratory oxygen fraction for preventing SSIs**

[Wetterslev 2015](#) included 28 trials (9330 participants in total). Of these, we categorised 15 included trials (7219 participants) as delivering interventions that started in the intraoperative phase.

### **SSI**

Fifteen relevant trials (7219 participants) reported SSI data, which we pooled. There was no clear difference in SSI risk following use of 60% to 90% oxygen compared with 30% to 40% oxygen (RR 0.87, 95% CI 0.71 to 1.07; low-certainty evidence; downgraded once for risk of bias and once for inconsistency - assessed by overview authors).

### **Mortality**

Eight relevant trials (4918 participants) in this review found no clear difference in mortality risk following use of 60% to 90% oxygen compared with 30% to 40% oxygen (RR 1.07, 95% CI 0.87 to 1.33; low-certainty evidence; downgraded once for imprecision and once for heterogeneity - assessed by overview authors). No other outcome data relevant to the overview were reported.

## **5. Use of antibiotics**

### **5.1. Antibiotic prophylaxis versus no prophylaxis for preventing infection after caesarean section**

[Smaill 2014](#) included 95 trials (more than 15,000 women). We categorised all the included trials as delivering interventions that

started in the intraoperative phase. In these trials antibiotic treatment was continued postoperatively.

### SSI

Eighty-two relevant trials (14,407 participants in total) reported SSI data presenting a single comparison of antibiotics with no antibiotics. Available trial evidence reports that antibiotic prophylaxis probably reduces SSIs (RR 0.40, 95% CI 0.35 to 0.46; moderate-certainty evidence; downgraded once for risk of bias - assessed by review authors).

No other outcome data relevant to the overview were reported.

### 5.2. Different classes of antibiotics given to women routinely for preventing SSI at caesarean section

Gyte 2014 included 31 RCTs (7697 participants in total).

### SSI

There were 19 relevant included trials (3559 participants in total). Of these 17 trials specified the timing of administration, which we categorised as the intraoperative phase, while two trials did not specify the timing of administration. These trials reported SSI data presenting four comparisons of different antibiotic prophylaxis regimes, including single cephalosporin, cephalosporin drug combination, single penicillin and penicillin drug combinations. All comparisons found no clear difference in SSI risk between the different regimes used (low- or very low-certainty evidence; variously downgraded once or twice for risk of bias and imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 5.3. Antibiotic prophylaxis for the prevention of methicillin-resistant *Staphylococcus aureus* (MRSA)-related complications in surgical patients

Gurusamy 2013 included 12 trials (4704 participants in total). Of these, we categorised seven trials (3393 participants) as delivering interventions that started in the intraoperative phase. All these trials continued antibiotic treatment postoperatively.

### SSI

Six trials (3294 participants in total) presented 11 comparisons of different prophylactic antibiotic regimens with each other, including pefloxacin, cefazolin, ertapenem, cefotetan, cefamandole, gentamycin, vancomycin, daptomycin, and cefuroxime. For all 11 comparisons there was no clear difference in SSI risk from use of one antibiotic prophylaxis regime compared with another (low- or very low-certainty evidence; variously downgraded for risk of bias and imprecision - assessed by overview authors). One trial

(99 participants) compared antibiotic prophylaxis (co-amoxiclav or cefotaxime) with no antibiotic prophylaxis and showed that receiving antibiotic prophylaxis with co-amoxiclav (or cefotaxime if allergic to penicillin) may reduce SSI risk (RR 0.26, 95% CI 0.11 to 0.65; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors).

### Mortality

Two relevant trials reported mortality. One trial (99 participants) found no clear difference in mortality risk when using co-amoxiclav or cefotaxime compared with no antibiotic prophylaxis (RR 0.54, 95% CI 0.17 to 1.72; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors). A second trial (884 participants) found no clear difference when using vancomycin compared with cefuroxime (RR 2.02, 95% CI 0.18 to 22.18; very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by overview authors). No other outcome data relevant to the overview were reported.

### 5.4. Prophylactic antibiotics for preventing SSIs after breast cancer surgery

Jones 2014 included 11 RCTs (2867 participants). Of these we categorised nine trials as delivering interventions that started in the intraoperative phase.

### SSI

Nine trials (2739 participants in total) presented three comparisons of different prophylactic antibiotic regimens. For the comparison of antibiotics delivered immediately prior to surgery compared with placebo, we pooled data from six trials (1708 participants): the use of antibiotic reduced SSI risk (RR 0.74, 95% CI 0.56 to 0.98; high-certainty evidence - assessed by overview authors). We also pooled the data from two trials (987 participants) and the use of antibiotics immediately prior to surgery may reduce the risk of SSIs compared with no treatment (RR 0.48, 95% CI 0.28 to 0.82; low-certainty evidence; downgraded once for imprecision and once for inconsistency - assessed by overview authors). One trial (44 participants) compared perioperative antibiotics with no antibiotic and found that it is uncertain whether perioperative antibiotics reduce SSI risk (very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 5.5. Systemic antimicrobial prophylaxis for preventing SSIs after percutaneous endoscopic gastrostomy (PEG)

Lipp 2013 included 13 RCTs (1637 participants in total) and we categorised all trials as delivering interventions that started in the intraoperative phase.

#### SSI

All trials reported peristomal infection as an outcome. A pooled analysis (by review authors) of 12 trials (1271 participants) found that prophylactic antibiotics may reduce the incidence of peristomal infection (RR 0.39, 95% CI 0.30 to 0.51; low-certainty evidence, downgraded twice for risk of bias - assessed by overview authors). Another trial (334 participants) compared intravenous (IV) antibiotics with antibiotics via PEG but the review authors could not include it in the meta-analysis. The evidence reported that it was uncertain whether there was a difference in peristomal infection risk following treatment with systemic antibiotic (PEG) compared with systemic antibiotic (IV) (RR 0.70, 95% CI 0.30 to 1.65 very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by overview authors). No other outcome data relevant to the overview were reported.

### 5.6. Timing of intravenous prophylactic antibiotics for preventing SSIs undergoing caesarean delivery

Mackeen 2014 included 10 trials (5041 participants in total) and we categorised all trials as delivering interventions that started in the intraoperative phase.

#### SSI

This review compared prophylactic intravenous antibiotics administered before caesarean incision with administration after cord clamping in a single comparison. Available trial evidence reports caesarean antibiotic prophylaxis administered intraoperatively prior to incision reduced maternal SSIs (RR 0.59, 95% CI 0.44 to 0.81; high-certainty evidence - assessed by review authors). No other outcome data relevant to the overview were reported.

### 5.7. Routes of administration of antibiotic prophylaxis for preventing infection after caesarean section

Nabhan 2016 included 10 trials (1354 participants in total). Of these we categorised seven trials (859 participants in total) as delivering interventions that started in the intraoperative phase.

#### SSI

Seven relevant trials (859 participants) reported SSI data. It is uncertain whether IV antibiotics reduce SSIs risk compared with irrigation (very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by review authors). No other outcome data relevant to the overview were reported.

### 5.8. Antibiotic prophylaxis for preventing SSIs in patients undergoing elective laparoscopic cholecystectomy

Sanabria 2010 included 11 trials (1664 participants in total). We categorised all included trials as delivering interventions that started in the intraoperative phase.

#### SSI

Eleven trials (1664 participants) reported SSI data presenting a single comparison of antibiotic prophylaxis with placebo or no prophylaxis. It is uncertain whether antibiotic prophylaxis reduces SSI risk in this comparison (very low-certainty evidence; downgraded twice for risk of bias and once for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 5.9. Antibiotic prophylaxis for hernia repair

Sanchez-Manuel 2012 included 17 trials (7843 participants in total). We categorised all included trials as delivering interventions that started in the intraoperative phase.

#### SSI

Seventeen trials (7843 participants) reported SSI data presenting a single comparison of antibiotic prophylaxis with placebo or no treatment. Available trial evidence reports that antibiotic prophylaxis probably reduces SSI risk (RR 0.67, 95% CI 0.54 to 0.84; moderate-certainty evidence; downgraded once for risk of bias - assessed by overview authors). Based on ROBIS, however, we assessed this review as being at unclear risk of bias due to a limited risk of bias assessment processes being used. This means that the overview authors were unable to fully assess the risk of bias for all domains recognised in the current version of the Cochrane 'Risk of bias' tool. We have not downgraded further for this review-level issue.

No other outcome data relevant to the overview were reported.

### 5.10. Antimicrobial prophylaxis for preventing SSIs in colorectal surgery

Nelson 2014 included 260 trials (43,451 participants in total) and 68 different antibiotics. Of these, we categorised 22 included trials

(3604 participants in total) as delivering interventions that started in the intraoperative phase.

## SSI

Twenty-two trials (3604 participants) presented six comparisons of different antibiotic regimens or different routes of administration of antibiotic prophylaxis. For the comparison of antibiotic with no antibiotic/placebo, we pooled the data from five trials (405 participants) and found that antibiotic may reduce SSI risk (RR 0.25, 95% CI 0.16 to 0.41; low-certainty evidence; downgraded once for risk of bias and once for imprecision - assessed by overview authors).

For the comparison of duration of therapy, we pooled data from seven trials (1484 participants) and found probably no difference in SSI risk with short-term compared with long-term duration antibiotic (RR 1.05, 95% CI 0.78 to 1.40; moderate certainty evidence; downgraded once for imprecision - assessed by overview authors).

For the comparison of additional aerobic coverage, we pooled data from four trials (230 participants) and found that, with added aerobic coverage, an antimicrobial prophylaxis regimen may slightly reduce SSI risk compared with no additional aerobic coverage (RR 0.38, 95% CI 0.16 to 0.96; low-certainty evidence; downgraded once for risk of bias and once for imprecision - assessed by overview authors).

For the comparison of additional anaerobic coverage, we pooled data from four trials (1098 participants) and found that, with added anaerobic coverage, an antimicrobial prophylaxis regimen may slightly reduce SSI risk compared with no additional anaerobic coverage (RR 0.65, 95% CI 0.47 to 0.90; low-certainty evidence; downgraded once for risk of bias and once for imprecision - assessed by overview authors).

For the comparisons of the different routes of administration of antibiotics from one trial (72 participants), it is uncertain whether oral antibiotics reduce SSI risk compared with intravenous routes (RR 2.11, 95% CI 0.20 to 22.29; very low-certainty evidence, downgraded once for risk of bias and twice for imprecision - assessed by overview authors). Evidence from one trial (310 participants) showed no clear difference when applying combined oral and intravenous antibiotics compared with oral or intravenous antibiotics alone (RR 0.50, 95% CI 0.23 to 1.11; low-certainty evidence, downgraded once for risk of bias and once for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 5.11. Methods of decreasing infection to improve outcomes after liver resections

[Gurusamy 2011](#) included seven trials (521 participants in total). Only two included trials reported mortality data, which we cate-

gorised as delivering interventions that started in the intraoperative phase.

## Mortality

One trial (180 participants) compared long-duration antibiotics with short-duration antibiotics; however there were no events in either arm in this trial (very low-certainty evidence; downgraded once for risk of bias, once for imprecision and once for publication bias - assessed by review authors). Another trial (59 participants) compared topical povidone iodine gel with no topical povidone iodine gel. It is uncertain whether topical povidone iodine gel reduces mortality risk (very low-certainty evidence; downgraded once for risk of bias, once for imprecision and once for publication bias - assessed by review authors).

No other outcome data relevant to the overview were reported.

### 5.12. Perioperative antibiotics to prevent infection after first-trimester abortion

[Low 2012](#) included 19 trials (9715 participants in total). No outcome data relevant to the overview were reported.

## 6. Management of theatre traffic

No reviews examined management of theatre traffic.

## 7. Wound irrigation

No reviews examined wound irrigation.

## 8. Wound closure

### 8.1. Continuous versus interrupted skin sutures for non-obstetric surgery

[Gurusamy 2014a](#) included five trials (827 participants in total) and of these, four trials (602 participants in total) reported SSI data.

## SSI

Evidence from four trials (602 participants) showed that it is uncertain whether continuous skin sutures reduce SSI risk compared with interrupted skin sutures (very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by review authors).

No other outcome data relevant to the overview were reported.

### 8.2. Subcutaneous closure versus no subcutaneous closure after non-caesarean surgical procedures

Gurusamy 2014b included six trials (815 participants in total) that compared subcutaneous closure with no subcutaneous closure in a single comparison.

#### SSI

Evidence showed that it is uncertain whether subcutaneous closure reduces SSI risk (very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by review authors). No other outcome data relevant to the overview were reported.

### 8.3. Techniques and materials for skin closure in caesarean section for preventing SSIs

Mackeen 2012 included seven trials (1104 participants in total) and presented two comparisons reporting SSI risk.

#### SSI

For the comparison of staples with absorbable subcuticular suture, data from six trials (916 participants) were pooled and there was no clear difference in SSI risk following use of absorbable subcuticular suture (RR 0.85, 95% CI 0.43 to 1.71; low-certainty evidence; downgraded once for risk of bias and once for inconsistency - assessed by overview authors). For the comparison of barbed suture with polydioxanone suture, data from one trial (188 participants) showed no clear difference in SSI risk when using the different types of sutures (RR 0.96, 95% CI 0.18 to 5.10; low-certainty evidence; downgraded twice for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 8.4. Healing by primary versus secondary intention after surgical treatment for pilonidal sinus

AL-Khamis 2010 included 26 trials (2530 participants in total) and of these, 17 trials (1940 participants) reported SSI data.

#### SSI

Data from 10 trials (1231 participants) showed that it is uncertain whether open healing reduces SSI risk compared with midline closure (RR 1.31, 95% CI 0.93 to 1.85; very low-certainty evidence; downgraded once for risk of bias, once for imprecision and once for inconsistency - assessed by overview authors). Data from five trials (541 participants) showed midline closure may increase the rate of SSIs compared with other closure (RR 3.72, 95% CI 1.86 to 7.42; low-certainty evidence; downgraded for risk of bias and

imprecision - assessed by overview authors). Evidence from one trial (68 participants) showed that it is uncertain whether classic Limberg reduces SSI risk compared with modified Limberg (very low-certainty evidence; downgraded once for risk of bias, twice for imprecision - assessed by overview authors). Similarly, evidence from another trial (100 participants) showed that it is uncertain whether classic Limberg reduces SSI risk compared with Karydakis (very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by overview authors).

No other outcome data relevant to the overview were reported.

### 8.5. Staples versus sutures for closing leg wounds after vein graft harvesting for coronary artery bypass surgery

Biancari 2010 included three trials (322 participants in total) that compared staple closure with suture closure in a single comparison.

#### SSI

It is uncertain whether staples reduce SSI risk compared with sutures (very low-certainty evidence; downgraded once for risk of bias and twice for imprecision - assessed by overview authors). No other outcome data relevant to the overview were reported.

### 8.6. Tissue adhesives for closure of surgical incisions

Dumville 2014 included 33 trials (2793 participants in total) and of these, 22 trials (1731 participants in total) reported SSI data.

#### SSI

Twenty-two trials (1731 participants in total) presented six comparisons of tissue adhesives with different wound-closing technologies. There was no clear difference in SSI risk between wounds closed with tissue adhesives and wounds closed using other methods reported (sutures, adhesive tape, staples or others) (low- or very low-certainty evidence; variously downgraded for risk of bias and imprecision - assessed by review authors).

No other outcome data relevant to the overview were reported.

### 9. Theatre cleansing

No reviews examined theatre cleansing.

## DISCUSSION

### Summary of main results



We have summarised the main results of the included reviews by categorising their findings and GRADE assessment ([GRADE 2013](#)) ([Table 7](#)).

The relative effects of majority of included interventions are inconclusive due to the low or very low certainty evidence. Exceptions to this are listed below. All data listed relate to SSI. There was no high or moderate certainty evidence for the relative effects of intra-operative interventions on mortality and no outcome data at all for quality of life or costs.

### High quality evidence

- Prophylactic intravenous antibiotics administered before caesarean incision reduce SSI risk compared with after neonatal umbilical cord clamping (RR 0.59, 95% CI 0.44 to 0.81; high-certainty evidence - rated by review authors).
- Adhesive drapes increase SSI risk compared with no drapes (RR 1.23, 95% CI 1.02 to 1.48; high-certainty evidence - rated by review authors) (negative effects).
- Preoperative antibiotics reduce SSI risk compared with placebo after breast cancer surgery (RR 0.74, 95% CI 0.56 to 0.98; high-certainty evidence - assessed by overview authors).

### Moderate quality evidence

- Antibiotic prophylaxis probably reduce SSI risk after caesarean section compared with no prophylaxis (RR 0.40, 95% CI 0.35 to 0.46; moderate-certainty evidence - rated by review authors).
- Antibiotic prophylaxis probably reduce SSI risk compared with placebo for hernia repair (RR 0.67, 95% CI 0.54 to 0.84; moderate-certainty evidence - rated by overview authors).
- Iodine-impregnated adhesive drapes (compared with no adhesive drapes); and duration of the use of antimicrobial prophylaxis for colorectal surgery (short-term compared with long-term duration antibiotic) probably lead to little difference in SSI risk (moderate-certainty evidence - rated by overview authors).

### Overall completeness and applicability of evidence

The evidence included in this overview covers all eligible Cochrane Reviews. Of the 32 included reviews, seven could be considered up-to-date as they were published within the last two years (Cochrane recommends updating reviews every two years) ([Campbell 2015](#); [Dumville 2015](#); [Nabhan 2016](#); [Vincent 2016](#); [Webster 2015](#); [Wetterslev 2015](#); [Wood 2016](#)).

In keeping with the nature of a Cochrane overview, this body of work does not cover non-Cochrane reviews. Alternative or emerging strategies for prevention of SSIs may not yet have been covered in a Cochrane Review and thus these data are not included

here, for example, use of Triclosan-containing sutures in children or laminar airflow ventilation systems. Once such strategies have been assessed in new reviews, we can and will update this overview accordingly.

### Quality of the evidence

In assessing the quality of the evidence, we employed the ROBIS tool to examine the reviews, and evaluated the authors' conclusions to ensure that they were appropriate based on the available data. All 32 included reviews scored well across the ROBIS assessment, likely due to the stringent reporting guidelines implemented by Cochrane prior to publication.

We used GRADE to assess the quality of the evidence reported by primary studies in the included reviews. The evidence presented in the majority of comparisons (91%) was rated either low- or very low-quality/certainty. The main reasons for downgrading the certainty of evidence included bias in the primary trials and imprecision, the latter caused by small sample sizes or low event rates, or both. It must be noted that the overview authors might have used different criteria to make GRADE assessments to the review authors. For example in our process we used an OIS (optimal information size) and this informed our decisions on downgrading for precision - this may not have been the case in other reviews. For transparency, we have reported review authors' GRADE decisions but these may not calibrate well with our assessments.

### Potential biases in the overview process

By only searching the Cochrane Library, and including only current Cochrane Reviews we may have missed some key literature. However, previous publications have referred to the higher-quality grading (high ROBIS score) in Cochrane Reviews due to the basic criteria necessary for publication at any stage (protocol or full review), suggesting that they may be the most reliable source of evidence ([Pollock 2017](#)).

We have employed a standard GRADE process on the included studies in reviews ([Schünemann 2011a](#); [Schünemann 2011b](#)). In one case we considered how a review-level issue of sub-optimal risk of bias assessment affected the GRADE assessment ([Sanchez-Manuel 2012](#)). In this case we did not alter the level of GRADE certainty given, but uncertainty on the quality of the review providing the evidence that was graded must be recognised.

### Agreements and disagreements with other studies or reviews

Over the years, as new evidence from RCTs continues to emerge, a steady stream of publications aim to provide a comprehensive overview on the prevention of SSIs. This is a summary overview

of current Cochrane Reviews, we are not aware of any similar overviews of prevention for SSIs.

World Health Organization (WHO) guidelines on SSI prevention have recently been published (WHO 2016). The WHO reviews that underpin these guidelines (WHO 2016) were also split by operative phase: preoperative, intraoperative and postoperative. The methods used to conduct the systematic reviews that underpin these guidelines were, in some cases, different to those of the corresponding Cochrane Reviews included in this overview of reviews, which means direct comparison between overview and guidelines findings is not appropriate. The WHO reviews are standard systematic reviews, more recent ones, in some cases, include observational as well as randomised controlled trial data and have review questions that, in some cases, differ in scope to corresponding Cochrane Reviews (as do related alibility criteria). Focusing on the respective findings of the guidelines and the overview for the intraoperative phase, the guidelines include topics not covered by Cochrane Reviews, such as maintenance of body temperature, maintenance of adequate circulating volume control, discontinuation of immunosuppressive agents and use of laminar airflow ventilation systems. Additionally, intraoperative antibiotic prophylaxis, considered as part of this overview, were only considered as pre- or postoperative interventions in the guidelines. Four further interventions were considered in both the guidelines and this overview: patient oxygenation, use of microbial sealants, blood glucose control and use of drapes. The WHO guidelines (WHO 2016) make a strong recommendation with moderate-quality evidence for use of 80% inspired oxygen intraoperatively and into the postoperative period for adult patients under general anaesthesia with endotracheal intubation. Our overview found low-certainty evidence from one review (Wetterslev 2015) with 15 RCTs reporting no clear difference in SSI risk following use of high perioperative inspiratory oxygen fraction for adult surgical patients. Although similar data were used in the analysis performed, unlike Wetterslev 2015, WHO 2016 conducted a subgroup analysis based on the type of anaesthesia, and it is this subgroup analysis that informs the recommendation made (on oxygenation). Use of surgical drapes was also considered by both guidelines and our overview. This overview considers two more RCTs than WHO guidelines, but both sources report similar findings in that adhesive drapes appear to increase the SSI risk compared with no drapes. Again, there were no key differences in findings reported for microbial sealant and blood glucose control.

## AUTHORS' CONCLUSIONS

### Implications for practice

This overview provides the most up-to-date evidence on prevention of SSIs from currently published Cochrane Reviews (intraop-

erative phase). Generally, we found insufficient or low-certainty evidence for the effect of most interventions for preventing SSIs. This comprehensive overview of Cochrane Reviews highlights the current uncertainty regarding the effectiveness of the intraoperative phase interventions as preventions for SSIs. It is important to note that one review with high-certainty evidence showed harms associated with the use of adhesive drapes; and another review also with high-certainty evidence showed benefit when using prophylactic intravenous antibiotics administered before caesarean incision. As there remains uncertainty on the use of a number of prophylactic SSI prevention options, health professionals are likely to follow local and national guidelines until more information becomes available.

### Implications for research

The individual reviews and this overview have highlighted the lack of good evidence for intraoperative interventions for SSI prevention. Included reviews in this category focused on interventions administered during the procedure (e.g. prophylactic antibiotics, patient warming) and methods to reduce bacterial contamination (e.g. glove changes, incise drapes). Just a few interventions altered the surgical approach itself (e.g. closure methods, the use of electrosurgical incisions). It is possible that different surgical techniques may influence SSI and this may be an area in need of more research. Most of the trials and the participants included in them did not contribute to any reliable assessment of efficacy or harm, which may lead to research waste. Robust randomised controlled trials with good internal validity from use of appropriate methods of randomisation, blinding and analysis are required. Studies also need to have carefully considered sample size calculations and recruitment strategies to ensure that they are not underpowered. It is also important that the outcomes that are important to patients and health professionals are measured. Future studies should use appropriate outcome measures that are consistent, reliable, have internal and external validity, and are sensitive to change in what is being measured. Consistent use of outcomes and related definitions would maximise the value of data from across multiple studies. Improving measurement of SSI, especially after hospital discharge, is warranted to improve data collection in this phase using validated patient-reported outcome (PRO) measures or methods for wound photography, or both, to complement these. A core outcome set focused on surgical wounds may be considered by developing and applying agreed, standardised sets of outcomes in this area. Trials should also collect quality-of-life data and consider incorporating cost-effectiveness analysis. Whilst adverse events should be collected as part of a trial, additional data on mortality and other rare events might be better collected as part of observational, prospective studies - perhaps using routinely collected data if possible. Crucially it is important to understand the risk of death as a function of SSI severity and these data are unlikely to be obtained from trials. This research also highlights the need for review authors to update existing reviews to ensure

that new studies are incorporated into existing reviews so that Cochrane Reviews remain contemporary and relevant.

## ACKNOWLEDGEMENTS

The authors would like to thank peer reviewers Kurinchi Gurusamy, Gill Worthy, Colin Davies and Janet Wale for their comments on the overview, and Lawrence Best and Ros Wade for their comments on the protocol. The reviewers are also grateful for copy-editors Ann Jones for editing the protocol and Denise Mitchell for editing the overview.

## REFERENCES

### References to included reviews

AL-Khamis A, McCallum I, King PM, Bruce J. Healing by primary versus secondary intention after surgical treatment for pilonidal sinus. *Cochrane Database of Systematic Reviews* 2010, Issue 1. [DOI: 10.1002/14651858.CD006213.pub3]

Biancari F, Tiozzo V. Staples versus sutures for closing leg wounds after vein graft harvesting for coronary artery bypass surgery. *Cochrane Database of Systematic Reviews* 2010, Issue 5. [DOI: 10.1002/14651858.CD008057.pub2]

Buchleitner AM, Martínez-Alonso M, Hernández M, Solà I, Mauricio D. Perioperative glycaemic control for diabetic patients undergoing surgery. *Cochrane Database of Systematic Reviews* 2012, Issue 9. [DOI: 10.1002/14651858.CD007315.pub2]

Campbell G, Alderson P, Smith AF, Wärttig S. Warming of intravenous and irrigation fluids for preventing inadvertent perioperative hypothermia. *Cochrane Database of Systematic Reviews* 2015, Issue 4. [DOI: 10.1002/14651858.CD009891.pub2]

Charoenkwan K, Iheozor-Ejirofor Z, Rerkasem K, Matovinovic E. Scalpel versus electrosurgery for major abdominal incisions. *Cochrane Database of Systematic Reviews* 2017, Issue 6. [DOI: 10.1002/14651858.CD005987.pub3]

Cook LA, Pun A, Gallo MF, Lopez LM, Van Vliet HA. Scalpel versus no-scalpel incision for vasectomy. *Cochrane Database of Systematic Reviews* 2014, Issue 3. [DOI: 10.1002/14651858.CD004112.pub4]

Dumville JC, Coulthard P, Worthington HV, Riley P, Patel N, Darcey J, et al. Tissue adhesives for closure of surgical incisions. *Cochrane Database of Systematic Reviews* 2014, Issue 11. [DOI: 10.1002/14651858.CD004287.pub4]

Dumville JC, McFarlane E, Edwards P, Lipp A, Holmes A, Liu Z. Preoperative skin antiseptics for preventing surgical wound infections after clean surgery. *Cochrane Database of Systematic Reviews* 2015, Issue 4. [DOI: 10.1002/14651858.CD003949.pub4]

Grocott MP, Dushianthan A, Hamilton MA, Mythen MG,

Harrison D, Rowan K, Optimisation Systematic Review Steering Group. Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery. *Cochrane Database of Systematic Reviews* 2012, Issue 11. [DOI: 10.1002/14651858.CD004082.pub5]

Gurusamy KS, Naik P, Davidson BR. Methods of decreasing infection to improve outcomes after liver resections. *Cochrane Database of Systematic Reviews* 2011, Issue 11. [DOI: 10.1002/14651858.CD006933.pub2]

Gurusamy KS, Koti R, Wilson P, Davidson BR. Antibiotic prophylaxis for the prevention of methicillin-resistant *Staphylococcus aureus* (MRSA) related complications in surgical patients. *Cochrane Database of Systematic Reviews* 2013, Issue 8. [DOI: 10.1002/14651858.CD010268.pub2]

Gurusamy KS, Toon CD, Allen VB, Davidson BR. Continuous versus interrupted skin sutures for non-obstetric surgery. *Cochrane Database of Systematic Reviews* 2014, Issue 2. [DOI: 10.1002/14651858.CD010365.pub2]

Gurusamy KS, Toon CD, Davidson BR. Subcutaneous closure versus no subcutaneous closure after non-caesarean surgical procedures. *Cochrane Database of Systematic Reviews* 2014, Issue 1. [DOI: 10.1002/14651858.CD010425.pub2]

Gyte GM, Dou L, Vazquez JC. Different classes of antibiotics given to women routinely for preventing infection at caesarean section. *Cochrane Database of Systematic Reviews* 2014, Issue 11. [DOI: 10.1002/14651858.CD008726.pub2]

Haas DM, Morgan S, Contreras K. Vaginal preparation with antiseptic solution before cesarean section for preventing postoperative infections. *Cochrane Database of Systematic Reviews* 2014, Issue 12. [DOI: 10.1002/14651858.CD007892.pub5]

Hadiati DR, Hakimi M, Nurdiani DS, Ota E. Skin preparation for preventing infection following caesarean section. *Cochrane Database of Systematic Reviews* 2014, Issue 9. [DOI: 10.1002/14651858.CD007462.pub3]

Jones DJ, Bunn F, Bell-Syer SV. Prophylactic antibiotics to prevent surgical site infection after breast cancer surgery. *Cochrane Database of Systematic Reviews* 2014, Issue 3.



[DOI: 10.1002/14651858.CD005360.pub4]  
 Kao LS, Meeks D, Moyer VA, Lally KP. Peri-operative glycaemic control regimens for preventing surgical site infections in adults. *Cochrane Database of Systematic Reviews* 2009, Issue 3. [DOI: 10.1002/14651858.CD006806.pub2]  
 Lipp A, Lusardi G. Systemic antimicrobial prophylaxis for percutaneous endoscopic gastrostomy. *Cochrane Database of Systematic Reviews* 2013, Issue 11. [DOI: 10.1002/14651858.CD005571.pub3]  
 Low N, Mueller M, Van Vliet HA, Kapp N. Perioperative antibiotics to prevent infection after first-trimester abortion. *Cochrane Database of Systematic Reviews* 2012, Issue 3. [DOI: 10.1002/14651858.CD005217.pub2]  
 Mackeen AD, Berghella V, Larsen ML. Techniques and materials for skin closure in caesarean section. *Cochrane Database of Systematic Reviews* 2012, Issue 11. [DOI: 10.1002/14651858.CD003577.pub3]  
 Mackeen AD, Packard RE, Ota E, Berghella V, Baxter JK. Timing of intravenous prophylactic antibiotics for preventing postpartum infectious morbidity in women undergoing cesarean delivery. *Cochrane Database of Systematic Reviews* 2014, Issue 12. [DOI: 10.1002/14651858.CD009516.pub2]  
 Nabhan AF, Allam NE, Abdel-Aziz Salama MH. Routes of administration of antibiotic prophylaxis for preventing infection after caesarean section. *Cochrane Database of Systematic Reviews* 2016, Issue 6. [DOI: 10.1002/14651858.CD011876.pub2]  
 Nelson RL, Gladman E, Barbateskovic M. Antimicrobial prophylaxis for colorectal surgery. *Cochrane Database of Systematic Reviews* 2014, Issue 5. [DOI: 10.1002/14651858.CD001181.pub4]  
 Sanabria A, Dominguez LC, Valdivieso E, Gomez G. Antibiotic prophylaxis for patients undergoing elective laparoscopic cholecystectomy. *Cochrane Database of Systematic Reviews* 2010, Issue 12. [DOI: 10.1002/14651858.CD005265.pub2]  
 Sanchez-Manuel FJ, Lozano-García J, Seco-Gil JL. Antibiotic prophylaxis for hernia repair. *Cochrane Database of Systematic Reviews* 2012, Issue 2. [DOI: 10.1002/14651858.CD003769.pub4]  
 Smaill FM, Grivell RM. Antibiotic prophylaxis versus no prophylaxis for preventing infection after cesarean section. *Cochrane Database of Systematic Reviews* 2014, Issue 10. [DOI: 10.1002/14651858.CD007482.pub3]  
 Tanner J, Parkinson H. Double gloving to reduce surgical cross-infection. *Cochrane Database of Systematic Reviews* 2006, Issue 3. [DOI: 10.1002/14651858.CD003087.pub2]  
 Vincent M, Edwards P. Disposable surgical face masks for preventing surgical wound infection in clean surgery. *Cochrane Database of Systematic Reviews* 2016, Issue 4. [DOI: 10.1002/14651858.CD002929.pub3]  
 Webster J, Alghamdi A. Use of plastic adhesive drapes during surgery for preventing surgical site infection. *Cochrane Database of Systematic Reviews* 2015, Issue 4. [DOI: 10.1002/14651858.CD006353.pub4]  
 Wetterslev J, Meyhoff CS, Jørgensen LN, Gluud

C, Lindschou J, Rasmussen LS. The effects of high perioperative inspiratory oxygen fraction for adult surgical patients. *Cochrane Database of Systematic Reviews* 2015, Issue 6. [DOI: 10.1002/14651858.CD008884.pub2]  
 Wood C, Phillips C. Cyanoacrylate microbial sealants for skin preparation prior to surgery. *Cochrane Database of Systematic Reviews* 2016, Issue 5. [DOI: 10.1002/14651858.CD008062.pub4]

## References to excluded reviews

Cirocchi R, Randolph JJ, Montedori A, Cochetti GG, Arezzo A, Mearini EE, et al. Staples versus sutures for surgical wound closure in adults. *Cochrane Database of Systematic Reviews* 2014, Issue 8. [DOI: 10.1002/14651858.CD011250]  
 McCallum I, O'Loughlin P, Robinson S, Wright LJ, Thomas CL. Antibiotic-coated sutures versus non-antibiotic-coated sutures for the prevention of abdominal surgical wound infection. *Cochrane Database of Systematic Reviews* 2016, Issue 11. [DOI: 10.1002/14651858.CD011297.pub2]  
 Ousey KJ, Edward KL, Lui S, Stephenson J, Duff J, Walker KN, et al. Perioperative warming therapy for preventing surgical site infection in adults undergoing surgery. *Cochrane Database of Systematic Reviews* 2016, Issue 12. [DOI: 10.1002/14651858.CD011731.pub2]  
 Smith TA, Rowlands C, Dumville JC, Norman G. Intracavity lavage and wound irrigation for prevention of surgical site infection. *Cochrane Database of Systematic Reviews* 2016, Issue 6. [DOI: 10.1002/14651858.CD012234]  
 Verschuur HP, De Wever W, Van Benthem PP. Antibiotic prophylaxis in clean and clean-contaminated ear surgery. *Cochrane Database of Systematic Reviews* 2004, Issue 3. [DOI: 10.1002/14651858.CD003996.pub2]

## Additional references

### ACORN 2012

Australian College of Operating Room Nurses (ACORN). *Standards for Perioperative Nursing*. 14th Edition. Adelaide, Australia: ACORN, 2012.

### Al-Niaimi 2009

Al-Niaimi A, Safdar N. Supplemental perioperative oxygen for reducing surgical site infection: a meta-analysis. *Journal of Evaluation in Clinical Practice* 2009;**15**(2):360–5.

### Allegranzi 2010

Allegranzi B, Nejad SB, Combescure C, Graafmans W, Attar H, Donaldson L, et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet* 2010;**377**(9761):228–41.

### Allegranzi 2016a

Allegranzi B, Bischoff P, de Jonge S, Kubilay NZ, Zayed B, Gomes SM, et al. WHO Guidelines Development Group. New WHO recommendations on preoperative measures for surgical site infection prevention: an evidence-based global perspective. *Lancet Infectious Diseases* 2016;**16**(12):e276–87.

**Allegranzi 2016b**

Allegranzi B, Zayed B, Bischoff P, Kubilay NZ, de Jonge S, de Vries F, et al. WHO Guidelines Development Group. New WHO recommendations on intraoperative and postoperative measures for surgical site infection prevention: an evidence-based global perspective. *Lancet Infectious Diseases* 2016;**16**(12):e288–303.

**ASA 2014**

American Society of Anesthesiologists (ASA). ASA physical status classification system. October 2014. [www.asahq.org/resources/clinical-information/asa-physical-status-classification-system](http://www.asahq.org/resources/clinical-information/asa-physical-status-classification-system) (accessed 4 October 2017).

**Astagneau 2009**

Astagneau P, L'Hériveau F, Daniel F, Parneix P, Venier AG, Malavaud S, et al. Reducing surgical site infection incidence through a network: results from the French ISO-RAISIN surveillance system. *Journal of Hospital Infection* 2009;**72**(2):127–34.

**Awad 2012**

Awad SS. Adherence to surgical care improvement project measures and post-operative surgical site infections. *Surgical Infections* 2012;**13**(2):234–7.

**Barnes 2014**

Barnes S, Spencer M, Graham D, Johnson HB. Surgical wound irrigation: a call for evidence-based standardization of practice. *American Journal of Infection Control* 2014;**42**(5):525–9.

**Becker 2011**

Becker LA, Oxman AD. Chapter 22: Overviews of reviews. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from [handbook.cochrane.org](http://handbook.cochrane.org).

**Brown 2014**

Brown B, Tanner J, Padley W. This wound has spoiled everything: emotional capital and the experience of surgical site infections. *Sociology of Health and Illness* 2014;**36**(8):1171–87.

**Bruce 2001**

Bruce J, Russell EM, Mollison J, Krukowski ZH. The measurement and monitoring of surgical adverse events. *Health Technology Assessment* 2001;**5**(22):1–194.

**CDC 2017**

US Centers for Disease Control and Prevention (CDC). Surgical site infection (SSI) event. Procedure-associated module. January 2017. [www.cdc.gov/nhsn/PDFs/pscmanual/9pscasicurrent.pdf](http://www.cdc.gov/nhsn/PDFs/pscmanual/9pscasicurrent.pdf) (accessed 20 January 2017).

**Cooper 2003**

Cooper BS, Stone SP, Kibbler CC, Cookson BD, Roberts JA, Medley GF, et al. Systematic review of isolation policies in the hospital management of methicillin-resistant *Staphylococcus aureus*: a review of the literature with epidemiological and economic modelling. *Health Technology Assessment* 2003;**7**(39):1–194.

**De Lissovoy 2009**

De Lissovoy G, Fraeman K, Hutchins V, Murphy D, Song D, Vaughn BB. Surgical site infection: incidence and impact on hospital utilization and treatment costs. *American Journal of Infection Control* 2009;**37**(5):387–97.

**ECDC 2013**

European Centre for Disease Prevention and Control (ECDC). Point prevalence survey of healthcare associated infections and antimicrobial use in European acute care hospitals 2011–2012. [ecdc.europa.eu/en/publications/publications/healthcare-associated-infections-antimicrobial-use-pps.pdf](http://ecdc.europa.eu/en/publications/publications/healthcare-associated-infections-antimicrobial-use-pps.pdf) (accessed 12 December 2016).

**Gaynes 2001**

Gaynes RP, Culver DH, Horan TC, Edwards JR, Richards C, Tolson JS. Surgical site infection (SSI) rates in the United States, 1992–1998: the National Nosocomial Infections Surveillance System basic SSI risk index. *Clinical Infectious Diseases* 2001;**33**(Suppl 2):S69–77.

**Gibbons 2011**

Gibbons C, Bruce J, Carpenter J, Wilson AP, Wilson J, Pearson A, et al. Identification of risk factors by systematic review and development of risk-adjusted models for surgical site infection. *Health Technology Assessment* 2011;**15**(30):1–156.

**Goodman 2017**

Goodman T, Spry C. Introduction to perioperative nursing. In: Goodman T, Spry C editor(s). *Essentials of Perioperative Nursing*. 6th Edition. Burlington, MA: Jones & Barlett Learning, 2017:1–14.

**GRADE 2013**

Schünemann H, Brożek J, Guyatt G, Oxman A, GRADE working group. GRADE handbook. October 2013. [gdt.guidelinedevelopment.org/central\\_prod/\\_design/client/handbook/handbook.html](http://gdt.guidelinedevelopment.org/central_prod/_design/client/handbook/handbook.html) (accessed 5 February 2016).

**Greif 1999**

Greif R, Laciny S, Rapf B, Hickie RS, Sessler DI. Supplemental oxygen reduces the incidence of postoperative nausea and vomiting. *Anesthesiology* 1999;**5**:1246–52.

**Guyatt 2011**

Guyatt GH, Oxman AD, Kunz R, Brożek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence: imprecision. *Journal of Clinical Epidemiology* 2011;**64**(12):1283–93.

**Hardin 1997**

Hardin W, Nichols R. Handwashing and patient skin preparation. In: Malangoni MA editor(s). *Critical Issues in Operating Room Management*. Philadelphia: Lippincott-Raven, 1997:133–49.

**Health Protection Agency 2015**

Health Protection Agency. Surveillance of surgical site infections in NHS hospitals in England 2014/2015. [www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/484874/Surveillance\\_of\\_Surgical\\_Site\\_Infections\\_in\\_NHS\\_Hospitals\\_in\\_England\\_report\\_2014-15.pdf](http://www.gov.uk/government/uploads/system/uploads/attachment_data/file/484874/Surveillance_of_Surgical_Site_Infections_in_NHS_Hospitals_in_England_report_2014-15.pdf) (accessed 12 December 2016).

**HICPAC 1999**

Hospital Infection Control Practices Advisory Committee (HICPAC). Guideline for prevention of surgical site infection 1999. *Infection Control and Hospital Epidemiology* 1999;**20**(4):259.

**Horan 1992**

Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori T. CDC definitions of nosocomial surgical site infections 1992: a modification of CDC definitions of surgical wound infections. *Infection Control and Hospital Epidemiology* 1992;**13**(10):606–8.

**Hospital Infection Society 2007**

Hospital Infection Society. Third prevalence survey of healthcare associated infections in acute hospitals in England 2006. [webarchive.nationalarchives.gov.uk/20081105143757/dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_078388](http://webarchive.nationalarchives.gov.uk/20081105143757/dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_078388) (accessed 12 December 2016).

**Jenks 2014**

Jenks PJ, Laurent M, McQuarry S, Watkins R. Clinical and economic burden of surgical site infection and predicted financial consequences of elimination of SSI from an English hospital. *Journal of Hospital Infection* 2014;**86**(1): 24–33.

**Kovavisarach 2002**

Kovavisarach E, Seedadee C. Randomised controlled trial of glove perforation in single and double-gloving methods in gynaecologic surgery. *Australia and New Zealand Journal of Obstetrics and Gynaecology* 2002;**42**(5):519–21.

**Laine 2004**

Laine T, Aarnio P. Glove perforation in orthopaedic and trauma surgery: a comparison between single, double indicator and double gloving with two regular gloves. *Journal of Bone and Joint Surgery* 2004;**86**(6):898–900.

**Larson 1995**

Larson E. APIC guideline for handwashing and hand antisepsis in health-care settings. *American Journal of Infection Control* 1995;**23**(4):251–69.

**Ljungqvist 2005**

Ljungqvist O, Nygren J, Soop M, Thorell A. Metabolic perioperative management: novel concepts. *Current Opinion in Critical Care* 2005;**11**(4):295–9.

**Magill 2014**

Magill SS, Edwards JR, Bamburg W, Beldavs ZG, Dumyati G, Kainer MA, et al. Multistate point-prevalence survey of health care-associated infections. *New England Journal of Medicine* 2014;**370**(13):1198–208.

**Mangram 1999**

Mangram A, Horan T, Pearson M, Silver L, Jarvis W. Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection 1999. *Infection Control and Hospital Epidemiology* 1999;**20**(4): 250–78.

**NICE 2008**

National Institute for Health and Care Excellence (NICE). Surgical site infections: prevention and treatment. Clinical

guideline [CG74]. October 2008. [www.nice.org.uk/guidance/cg74](http://www.nice.org.uk/guidance/cg74) (accessed 23 February 2017).

**NICE 2016**

National Institute for Health and Care Excellence (NICE). Hypothermia: prevention and management in adults having surgery. Clinical guideline [CG65]. December 2016. [www.nice.org.uk/guidance/cg65](http://www.nice.org.uk/guidance/cg65) (accessed 23 February 2017).

**Pollock 2017**

Pollock A, Campbell P, Brunton G, Hunt H, Estcourt L. Selecting and implementing overview methods: implications from five exemplar overviews. *Systematic Reviews* 2017;**6**:145.

**Public Health England 2014**

Public Health England. Surveillance of surgical site infections in NHS hospitals in England: 2013 to 2014. [www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/386927/SSI\\_report\\_2013\\_14\\_final\\_3.pdf](http://www.gov.uk/government/uploads/system/uploads/attachment_data/file/386927/SSI_report_2013_14_final_3.pdf) (accessed 1 December 2016).

**Reichman 2009**

Reichman DE, Greenberg JA. Reducing surgical site infections: a review. *Reviews in Obstetrics and Gynecology* 2009;**2**(4):212–21.

**RevMan 2014 [Computer program]**

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

**Schünemann 2011a**

Schünemann HJ, Oxman AD, Higgins JP, Deeks JJ, Glasziou P, Guyatt GH. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from [handbook.cochrane.org](http://handbook.cochrane.org).

**Schünemann 2011b**

Schünemann HJ, Oxman AD, Higgins JP, Vist GE, Glasziou P, Guyatt GH. Chapter 11: Presenting results and 'Summary of findings' tables. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from [handbook.cochrane.org](http://handbook.cochrane.org).

**Singer 2008**

Singer AJ, Quinn JV, Hollander JE. The cyanoacrylate topical skin adhesives. *American Journal of Emergency Medicine* 2008;**26**(4):490–6.

**Smyth 2008**

Smyth ET, McIlvenny G, Enstone JE, Emmerson AM, Humphreys H, Fitzpatrick F, et al. Four country healthcare associated infection prevalence survey 2006: overview of the results. *Journal of Hospital Infection* 2008;**69**(3):230–48.

**Soderstrom 2003**

Soderstrom R. Principles of electrosurgery as applied to gynaecology. In: Rock JA, Jones HW III editor(s). *Te Linde's Operative Gynecology*. 9th Edition. Philadelphia: Lippincott Williams & Wilkins, 2003:291–308.

**Spagnolo 2013**

Spagnolo AM, Ottria G, Amicizia D, Perdellif, Cristina ML. Operating theatre quality and prevention of surgical site infections. *Journal of Preventive Medicine and Hygiene* 2013;**54**(3):131–7.

**Stephan 2002**

Stephan F, Yang K, Tankovic J, Soussy CJ, Dhonneur G, Duvaldestin P, et al. Impairment of polymorphonuclear neutrophil functions precedes nosocomial infections in critically ill patients. *Critical Care Medicine* 2002;**30**(2): 315–22.

**Swenson 2008**

Swenson BR, Camp TR, Mulloy DP, Sawyer RG. Antimicrobial-impregnated surgical incise drapes in the prevention of mesh infection after ventral hernia repair. *Surgical Infections* 2008;**9**(1):23–32.

**SWI Task Force 1992**

Surgical Wound Infection (SWI) Task Force. Consensus paper on the surveillance of surgical wound infections. *Infection Control and Hospital Epidemiology* 1992;**13**(10): 599–605.

**Wade 1980**

Wade A. *Pharmaceutical Handbook*. 19th Edition. London: Pharmaceutical Press, 1980.

**Warner 1988**

Warner C. Skin preparation in the surgical patient. *Journal of the National Medical Association* 1988;**80**(8):899–904.

**Whiting 2016**

Whiting P, Savović J, Higgins J, Caldwell D, Reeves B, Shea B, et al. ROBIS: a new tool to assess risk of bias in systematic reviews was developed. *Journal of Clinical Epidemiology* 2016; Vol. 69:225–34.

**Whitney 2015**

Whitney JD, Dellinger EP, Weber J, Swenson RE, Kent CD, Swanson PE, et al. The effects of local warming on surgical site infection. *Surgical Infections* 2015;**16**(5):595–603.

**WHO 2016**

World Health Organization (WHO). Global guidelines for the prevention of surgical site infection. November 2016. [www.who.int/gpsc/ssi-prevention-guidelines/en/](http://www.who.int/gpsc/ssi-prevention-guidelines/en/) (accessed 20 May 2017).

**References to other published versions of this review****Dumville 2017**

Dumville JC, Norman G, Westby MJ, Blazeby J, McFarlane E, Welton NJ, et al. Intra-operative interventions for preventing surgical site infection: an overview of Cochrane reviews. *Cochrane Database of Systematic Reviews* 2017, Issue 5. [DOI: 10.1002/14651858.CD012653

\* Indicates the major publication for the study

**ADDITIONAL TABLES****Table 1. Interventions aimed at preventing surgical site infections**

Intraoperative intervention types	Details	Theories on how the intervention type might work
For the patient		
Decontamination of patients' skin at site of surgery incision	Before surgery, patients' skin is disinfected using antiseptic solutions such as povidone-iodine or chlorhexidine at varying concentrations	The aim of preoperative skin antisepsis is to reduce the risk of SSIs by reducing the number of micro-organisms on the skin (ACORN 2012; Mangram 1999).
Skin sealants	Microbial sealants are liquids that are applied to the patient's skin before surgery and left to dry forming a protective film over the planned incision site. Cyanoacrylate, which is also used as a tissue adhesive, can be used as a skin sealant	As with other barrier methods, the use of skin sealants is focused on preventing contamination of the surgical wound with micro-organisms from the patient's skin. It is proposed that skin sealant use before surgery prevents any remaining micro-organisms from migrating into the surgi-

**Table 1. Interventions aimed at preventing surgical site infections** (Continued)

		cal wound following skin decontamination (Singer 2008).
Incise drapes	Before a surgical incision is made, sterile plastic adhesive (incise) drapes can be placed onto cleansed skin. The surgical incision is then made through the drape. Drapes can be plain or impregnated with antimicrobial products	Drapes are used as a barrier between the incision and the patient's skin, which, although cleansed may harbour micro-organisms, such as at deeper levels of the skin that cleansing cannot reach (Swenson 2008).
Use of electrosurgery for surgical incisions	In electrosurgery, an electric current is used to generate heat, which vaporises cellular material, cutting the skin in place of a scalpel. This can be used to cut skin from the top surface down or used on deep skin layers once an incision has been made with a scalpel (Soderstrom 2003).	It has been suggested that using heat to make a surgical incision may reduce the risk of SSI
Maintaining patient homoeostasis (warming)	During surgery the patient's bodily functions need to be optimised to promote recovery; it is further postulated this may also reduce the risk of SSI. Under general anaesthetic it is harder for the body to regulate its own temperature and this can increase the risk of perioperative hypothermia. Warming can be achieved using thermal insulation such as blankets, or active methods of warming that use machines to transfer heat to the patient, and use of heated intravenous fluids (NICE 2016; Whitney 2015).	Undertaking warming aims to maintain body temperature and prevent the development of perioperative hypothermia, which can lead to negative postoperative outcomes, which potentially include SSI. These interventions can also be used post-operatively to mitigate the impact of perioperative hypothermia when it has not been prevented
Maintaining patient homoeostasis (oxygenation)	During surgery under general anaesthetic patients are intubated and supplied with oxygen to maintain adequate oxygen perfusion to all tissues	It is suggested that the risk of SSI is higher when tissue oxygenation is not optimised during surgery. Some surgical protocols use higher saturation levels of oxygen during intubation to increase tissue oxygenation levels with the aim of reducing wound complications such as SSI. High oxygen levels have been linked to serious adverse events such as blindness and death (Al-Niaimi 2009).
Maintaining patient homoeostasis (blood glucose control)	Use of strict glycaemic control using medications to maintain glucose levels during surgery	Hyperglycaemia after surgery is postulated to lead to increased risk of surgical complications including infection (Ljungqvist 2005; Stephan 2002).

**Table 1. Interventions aimed at preventing surgical site infections** (Continued)

Wound irrigation and intracavity lavage (including use of intraoperative topical antiseptics before wound closure)	Surgical irrigation and intracavity lavage use fluids to wash out the surgical cavity at the end of the surgical procedure before the wound is closed. Both wound irrigation and intracavity lavage can be altered by: volume of irrigation fluid; mechanism or timing of delivery; or solution composition (Barnes 2014).	The theoretical advantage of surgical wound irrigation is to reduce the bacterial load in a surgical wound, and thus the risk of SSI, through a combination of water pressure, dilution, or the application of antimicrobial agents
Closure methods	Surgical wounds can be closed using sutures (absorbable or not) staples, adhesive strips or tissue adhesives. Some closure methods can make use of sutures that are coated in antimicrobial products The timing of closure can also vary; some wounds can be left open for a period following surgery and then closed (delayed closure)	There is a view that the method of surgical wound closure may impact on SSI risk. There is limited background evidence on mechanisms for SSI prevention, although it has been suggested that the better the seal the closure method obtains, the better the barrier to microbial contamination (Gurusamy 2014a).
For staff		
Use of masks, hair covers, overshoes, gowns and other protective coverings for theatre staff	Protective coverings worn in theatre by staff to limit the movement of micro-organisms in theatre (Cooper 2003). For example: masks over the face; disposable shoe covers worn over standard footwear and changed as required; disposable or reusable gowns worn over standard scrub outfits and changed as required	There are various coverings used in surgery that are designed to act as a barrier between the environment and the patient's wound to maintain a sterile operative field, such as masks that aim to capture water droplets being expelled. Masks contain one or two very finely woven filters that can inhibit bacteria. Masks cover the nose and mouth, but there is concern that masks may be worn incorrectly and allow air leaks from the sides of the mask Shoe coverings aim to limit the transfer of external material in and out of theatres Gowns cover standard surgical attire and can be removed when contaminated and replaced
Different glove protocols	Surgical staff wear disposable gloves during surgery. Gloves are used in a number of ways intended to minimise microbial contamination from staff to patients, including double gloving (using two pairs of gloves), the use of glove liners or cloth outer gloves (Kovavisarach 2002; Laine 2004).	Gloves are a barrier intervention that aim to prevent transfer of micro-organisms from the staff member's skin to the patient's skin or wound. Gloves also act as a barrier to prevent staff from infection by patients
For the environment		

**Table 1. Interventions aimed at preventing surgical site infections** (Continued)

Theatre cleansing	<p>The theatre environment needs to be cleaned regularly with detergents to disinfect surfaces. Daily deep cleaning is likely to occur using various protocols for cleaning surfaces between patient surgeries, especially areas that are contaminated with bodily fluid, or that are frequently touched by staff. Recent technologies used for theatre cleansing include UVC light decontamination and hydrogen peroxide vapour treatment</p> <p>Surgical instruments are also sterilised to decontaminate them after use. Various protocols are used including steam sterilisation and chemical sterilisation, which is used when steam sterilisation is not feasible</p> <p>Theatre cleaning can also involve the use of ventilation systems, such as laminar airflow systems, which supply filtered air into the environment to limit numbers of airborne micro-organisms</p> <p>To avoid cross-infection, special protocols may be developed for cleansing when surgical patients are known to have specific infections</p>	All aspects of theatre cleansing aim to minimise numbers of micro-organisms present in the theatre environment with the aim of reducing the risk of SSI. ( <a href="#">Spagnolo 2013</a> ).
Theatre traffic	A surgical theatre can be a busy working environment with people moving in and out. This movement can be managed, for example limiting the entrance and exit of staff during surgery, and minimising visitors into the theatre (e.g. partners of women undergoing caesarean sections) ( <a href="#">Spagnolo 2013</a> ).	A key aim in the prevention of SSI is to limit numbers of micro-organisms in the operative environment. People moving in and out of the operative field may increase the risk of contamination. Visitors to the theatre who have not undergone full hand scrubbing protocols and so forth could also potentially increase SSI risk

Table 2. Characteristics of included reviews

Review no.	First review author + year	Review title	Total number of included RCTs (and participants)	Review objective	Population, including types of surgery/ procedure and depth of incision	Main intervention e. g. nasal decontamination	Comparator (s)	Relevant review outcomes		Review limitations	Note
								Primary	Secondary		
CD006213	AL-Khamis 2010	Healing by primary versus secondary intention after surgical treatment for pilonidal sinus	26 studies (n = 2530)	To determine the relative effects of open compared with closed surgical treatment for pilonidal sinus on the outcomes of time to healing, infection and recurrence rate	Any participants (over 14 years of age) undergoing surgery to treat pilonidal sinus disease; surgical treatment for pilonidal sinus; variations of depth of incision (no details)	Any surgical intervention where the wound was left open to heal or closed by sutures	Another surgical intervention	Time to healing SSI Recurrence	Time to return to work Other complications and morbidity Participant (patient) satisfaction Cost Length of hospital stay Pain Quality of life Rate of change of wound volume Wound healing rate Operative time	Variations in the surgical techniques included in each group when conducting meta-analysis	They also compared different closed surgical treatments (midline vs off-midline wound closure Within each group there were variations in the surgical techniques used: for example, the amount of tissue excised, depth of incision,



**Table 2. Characteristics of included reviews** (Continued)

											type of suture
CD00805	<a href="#">Biancari 2010</a>	Staples versus sutures for closing leg wounds after vein graft harvesting for coronary artery bypass surgery	4 studies (n = 839 leg wounds in 581 participants)	To compare the rates of SSI and wound dehiscence of staples and sutures for skin closure after saphenous vein graft harvesting for coronary artery bypass graft surgery	People undergoing saphenous vein graft harvesting for CABG; minimally invasive vein harvesting was excluded	Suture	Staples	Rates of SSI Severity of SSI Time to wound healing	Rate of wound dehiscence Length of hospital stay Pain Cost Patient comfort Lower limb revascularization		Only 3 studies included (322 legs) were pooled into meta-analysis 1 study was excluded from the pooled analysis because each wound experienced both methods of closure and there was the risk of a unit of analysis error. However there was no statistically significant difference between the groups in this study.

**Table 2. Characteristics of included reviews** (Continued)

CD00731	<a href="#">Buch-leitner 2012<sup>a</sup></a>	Periop-er-ative gly-caemic control for dia-betic pa-tients un-der-go-ing surgery	12 stud-ies (n = 1403)	To assess the effects of periop-er-ative gly-caemic control for peo-ple with diabetes un-der-go-ing surgery	Partici-pants of any age, sex or ethnicity with previ-ously di-agnosed type 1 or 2 diabetes mellitus and submit- ted to periop-erative gly-caemic control	Periop-er-ative gly-caemic control proto- col pro- posed by study authors that in- volves a more in- ten- sive con- trol than the con- ven- tional care	Periop-er-ative gly-caemic con- trol pro- tocol de- fined as stan- dard or con- ven- tional care by the study authors	Any kind of infec- tious compli- cation All- cause mortal- ity Hypo- gly- caemic episodes	Cardio-vascular events Renal failure Length of ICU and hos- pital stay Health- related quality of life Eco- nomical costs Weight gain Mean blood glucose during inter- vention		
CD00989	<a href="#">Camp- bell 2015</a>	Warm- ing of in- tra- venous and irri- gation fluids for prevent- ing inad- ver- tent pe- rioper- ative hy- pother- mia	24 stud-ies (n = 1250)	To estimate the ef- fective- ness of preoper- ative or intraop- erative warm- ing, or both, of in- tra- venous and irri- gation fluids in pre- venting periop- erative hy- pother-	Adults un- der-go- ing elective or emer- gency surgery (in- cluding surgery for trauma) un- der general or re- gional (central neu- raxial block) anaes- thesia,	Warmed in- tra- venous fluids in- cluding all meth- ods of warm- ing flu- ids be- fore ad- minis- tration to the patient Warmed irri- ga- tion fluids in- cluding any irri- gation	Other warmed fluid inter- ventions Stan- dard care Thermal insula- tion or passive warm- ing Active warm- ing Preoper- ative or intraop- erative warm-	Risk of hy- pother- mia at any point during surgery and temper- ature at the end of surgery or on admis- sion to postanae- sthesia care; Major	Infec- tion and compli- cations of the sur- gical wound Pressure ulcers Bleeding compli- cations Other cardio- vascular com- plication Patient- reported out-		No data of inter- est to overview authors reported

**Table 2. Characteristics of included reviews** (Continued)

				mia and its complications during surgery in adults	or both	fluids administered to a body cavity that is warmed by any method	ing, or both, of inspired and insufflated gases Preoperative and intraoperative pharmacological interventions	cardio-vascular complications	comes All-cause mortality Length of stay Unplanned high dependency or intensive care admission Adverse effects		
CD00598	<a href="#">Charoenk 2017</a>	Scalpel versus electro-surgery for major abdominal incisions	16 studies (n = 2769)	To assess the effects of electro-surgery compared with scalpel for major abdominal incisions	People undergoing major open abdominal surgery, regardless of the orientation of the incision (vertical, oblique, or transverse) and surgical setting (elective or emergency)	Wound creation using electro-surgery	Wound creation using a scalpel	Wound infection Time to wound healing Wound dehiscence	Wound incision time Wound-related blood loss Postoperative pain Adhesion or scar formation		Sub-group analysis was planned but not possible to carry out due to interventions being insufficiently homogeneous and badly reported. Sensitivity analysis was planned by excluding stud-

**Table 2. Characteristics of included reviews** (Continued)

										ies at high or unclear risk of bias. However, this was not possible as none of the included studies were at low risk of bias
CD00411	<a href="#">Cook 2014</a>	Scalpel versus no-scalpel incision for vasectomy	2 studies (n = 1529)	To compare the effectiveness, safety, and acceptability of the incisional versus no-scalpel approach to the vasectomy	Men of reproductive age undergoing vasectomy for sterilisation	No-scalpel	Scalpel	Post-vasectomy adverse events (including wound infection)	Operating time Pain Time to resumption of intercourse Rates for azoospermia Time to azoospermia Pregnancy Incidence of recanalization Incidence of repeat vasectomy Cost analysis Consumer	

**Table 2. Characteristics of included reviews** (Continued)

									accept- abil- ity mea- sures Provider accept- abil- ity mea- sures		
CD00428	<a href="#">Dumville 2014</a>	Tissue adhe- sives for closure of surgi- cal inci- sions	33 stud- ies (n = 2793)	To de- termine the effects of various tissue adhe- sives com- pared with conven- tional skin closure tech- niques for the closure of sur- gical wounds	People of any age and in any set- ting re- quir- ing clo- sure of a surgical skin in- cision of any length	Tissue adhesive	An- other tis- sue ad- hesive or alterna- tive con- ven- tional closure device	Wound dehis- cence	Propor- tion of infected wounds Cos- metic appear- ance Pa- tient sat- isfaction Sur- geon sat- isfaction Cost Time taken to wound closure		
CD00394	<a href="#">Dumville 2015</a>	Preoper- ative skin an- tiseptics for pre- venting surgical wound infec- tions af- ter clean surgery	13 stud- ies (n = 2623)	To de- termine whether preop- erative skin an- tiseptics imme- diately prior to surgical incision for clean surgery prevents SSI and	People of any age un- dergo- ing clean surgery	Antisep- tic solu- tions or powders	A con- trol; an- other type of anti- septic or differ- ent dose	SSI	Quality of life Adverse events Re- source use		

**Table 2. Characteristics of included reviews** (Continued)

				to determine the comparative effectiveness of alternative antiseptics							
CD00408	<a href="#">Grocott 2012<sup>a</sup></a>	Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery	31 studies (n = 5292)	To describe the effects of increasing perioperative blood flow using fluids with or without inotropes or vasoactive drugs. Outcomes were mortality, morbidity, resource utilization and health status	Adults (aged $\geq$ 16 years) undergoing surgery in an operating theatre	Perioperative administration (initiated within 24 h before surgery and lasting up to 6 h after surgery) of fluids, with or without inotropes or vasoactive drugs to increase global blood flow against explicit measured goals	Control	Mortality (at longest available follow-up)	Mortality: all reported time frames Morbidity Resource utilisation Health status		Sub-group analysis and sensitivity analysis were done
CD00872	<a href="#">Gyte 2014<sup>a</sup></a>	Different classes of antibiotics	31 studies (n = 7697 women)	To determine, from the best available	Women undergoing caesarean	Prophylactic antibiotic regimens	Different classes of antibiotics ( $\geq$	Maternal: maternal sepsis (sus-	Maternal: fever (febrile morbid-		Sub-group analyses were carried out

**Table 2. Characteristics of included reviews** (Continued)

		given to women routinely for preventing infection at caesarean section	35 studies included in the review but only 31 provided data	evidence, the balance of benefits and harms between different classes of antibiotic given prophylactically to women undergoing caesarean section	section, both elective and non-elective		2 antibiotics from the different classes of antibiotics)	pected or proven); endometritis Infant: infant sepsis (suspected or proven); oral thrush	ity) ; wound infection; urinary tract infection; thrush; serious infectious complication; adverse effects of treatment on the woman; maternal lengths of hospital stay; infections - post-hospital discharge to 30 days postoperatively; readmissions Infant: immediate adverse effects of antibiotics on		by type of surgery; by time of administration; by route of administration Sensitivity analysis was not performed
--	--	--	---	--	---	--	--	--	---	--	--

**Table 2. Characteristics of included reviews** (Continued)

									the infant; infant length of hospital stay; long-term adverse effects; infant's immune system development Additional outcomes: development of bacterial resistance; costs		
CD00693	<a href="#">Gurusamy 2011<sup>a</sup></a>	Methods of decreasing infection to improve outcomes after liver resections	7 studies (n = 521)	To determine the benefits and harms of different interventions in decreasing the infectious complications and improving the outcomes after liver resection	People undergoing liver resection	Antibiotics Prebiotics or probiotics Immunomodulation Topical antibiotic or antiseptic	No antibiotics or placebo; no probiotics or probiotics or placebo; no immunomodulation; no topical antibiotic or antiseptic or saline or placebo before	Mortality Serious adverse events Quality of life	Hospital stay Number of unplanned visits to the doctor Return to work Costs		The unit of analysis was the aggregate data on participants undergoing liver resection according to randomised group Sensitivity analysis



**Table 2. Characteristics of included reviews** (Continued)

							wound closure; another of the included interventions				and subgroup analysis were not conducted
CD01026	<a href="#">Gurusamy 2013<sup>a</sup></a>	Antibiotic prophylaxis for the prevention of methicillin-resistant Staphylococcus aureus (MRSA) related complications in surgical patients	12 studies (n = 4704)	To compare the benefits and harms of all methods of antibiotic prophylaxis in the prevention of postoperative MRSA infection and related complications in people undergoing surgery	People undergoing surgery, irrespective of age, type of surgery, whether surgery was elective or emergency, and whether MRSA colonisation was identified by routine screening	Antibiotic prophylaxis	Placebo (or no treatment) ; different antibiotic prophylaxis (and regimens)	All-cause mortality Other serious adverse events Quality of life	Total length of hospital stay Use of health care resources Rates of SSIs Rates of SSIs due to MRSA Rates of infections due to MRSA		No subgroup analysis performed. Sensitivity analysis was done
CD01036	<a href="#">Gurusamy 2014a</a>	Continuous versus interrupted skin sutures for non-obstetric surgery	5 studies (n = 827)	To compare the benefits and harms of continuous compared with interrupted skin closure	People, of any age and sex, undergoing non-obstetric surgery	Continuous sutures	Interrupted sutures	SSI Wound dehiscence Quality of life	Hyper-trophic scarring Keloid scarring Incisional hernia Hospital stay Impact to the pa-		No subgroup analysis performed; sensitivity analysis was done

**Table 2. Characteristics of included reviews** (Continued)

				techniques in participants undergoing non-obstetric surgery					tient and to the health-care funder		
CD01042	<a href="#">Gurusamy 2014b</a>	Subcutaneous closure versus no subcutaneous closure after non-caesarean surgical procedures	6 studies (n = 815) 8 studies (n = 1318) included in the review but only 6 contributed data	To compare the benefits (such as decreased wound-related complications) and consequences (such as increased operating time) of subcutaneous closure compared with no subcutaneous closure in participants undergoing non-caesarean surgical procedures	People, of any age and sex, undergoing non-caesarean surgery	Subcutaneous closure	No subcutaneous closure, irrespective of the suture material	SSI Wound dehiscence Quality of life	Hyper-trophic scarring Keloid scarring Incisional hernia Hospital stay Impact to the patient and to the health-care funder		No subgroup analysis performed; sensitivity analysis was done

**Table 2. Characteristics of included reviews** (Continued)

CD00789	<a href="#">Haas 2014</a>	Vaginal preparation with antiseptic solution before cesarean section for preventing post-operative infections	7 studies (n = 2816; 2635 analysed)	To determine if cleansing the vagina with an anti-septic solution before a caesarean delivery decreases the risk of maternal infectious morbidities, including endometritis and wound complications	Pregnant women who received a caesarean delivery	Vaginal cleansing with any type of antiseptic solution	Placebo solution/standard care	Postpartum endometritis	Wound infection; fever; wound seroma or hematoma Composite wound complications Side effects of vaginal preparation		Sub-group analysis was done
CD00746	<a href="#">Hadiati 2014</a>	Skin preparation for preventing infection following caesarean section	6 studies (n = 1522)	To compare the effects of different agent forms and methods of pre-operative skin preparation for preventing post caesarean infection	Pregnant women undergoing elective or emergency caesarean section	Antiseptic agents used for caesarean section skin preparation	Different antiseptic agents, forms or methods of application	SSI Metritis or endometritis	Length of stay Maternal mortality Repeat surgery Re-admission resulting from infection Reduction of skin bacteria colony		No sub-group analysis performed; sensitivity analysis was not done

**Table 2. Characteristics of included reviews** (Continued)

				tion					count Adverse events		
CD00536	<a href="#">Jones 2014<sup>a</sup></a>	Prophy-lactic antibiotics to prevent sur-gical site infec-tion af-ter breast cancer surgery	11 stud-ies (n = 2867)	To de-terminate the effects of prophylactic (pre- or perioperative) antibiotics on the inci-dence of surgical site infection (SSI) after breast cancer surgery	People with breast cancer under-going breast surgery with or without im-me-diate recon-struction as part of their treat-ment	Any pre- or peri-operative anti-biotic used as prophylaxis where there was no known infec-tion	No an-tibiotic; placebo; another anti-biotic only if there was a control or placebo arm	SSI Ad-verse re-actions	Death Delay in adjuvant cancer treat-ment be-cause of breast wound infec-tion Time to wound healing Time to infec-tion Read-mission to hospi-tal Cost of care (should be a compar-ison be-tween the treat-ment and con-trol group)		Sensitiv-ity anal-ysis was per-formed
CD00680	<a href="#">Kao 2009<sup>a</sup></a>	Peri-operative gly-caemic con-trol regi-mens for preventing sur-gical site	5 studies (n = 773)	To sum-marise the evidence for the impact of gly-caemic control in the	People aged $\geq 18$ years , regard-less of diabetes status, who un-derwent a surgi-	1 gly-caemic control regimen	At least 1 other gly-caemic con-trol regi-men pre-, intra-, and/or postop-	SSI	Incidence and severity of hy-pogly-caemia Level of gly-		Sub-group analysis was per-formed (people with and with-out dia-

**Table 2. Characteristics of included reviews** (Continued)

		infections in adults		perioperative period on the incidence of surgical site infections, hypoglycaemia, level of glycaemic control, all-cause and infection-related mortality, and hospital length of stay and to investigate for differences of effect between different levels of glycaemic control	cal procedure		eratively		caemic control All-cause and infection-related mortality Length of hospital stay		betes); sensitivity analyses were not undertaken
CD00557	<a href="#">Lipp 2013<sup>a</sup></a>	Systemic antimicrobial prophylaxis for percutaneous endoscopic gastrostomy	13 studies (n = 1637)	To establish whether prophylactic use of systemic antimicrobials reduces the risk of peristomal	People of any age, gender or diagnosis, undergoing placement of a PEG tube	Antimicrobial prophylaxis	Placebo or usual care and comparisons between different antimicrobial regimens	Peristomal site infection	Identification of bacteria causing infection Peritonitis Adverse effects Mortality Re-		

**Table 2. Characteristics of included reviews** (Continued)

				infection in people undergoing placement of percutaneous endoscopic gastrostomy tubes					removal of PEG tube because of infection Length of hospital stay		
CD00521	Low 2012 <sup>a</sup>	Perioperative antibiotics to prevent infection after first-trimester abortion	19 studies (n = 9715)	To determine: 1. the effectiveness of antibiotic prophylaxis in preventing post-abortion upper genital tract infection; 2. the most effective antibiotic regimen; 3. the most effective strategy	All women undergoing induced first trimester surgical or medical abortion with or without a history of pelvic inflammatory disease, or a pre-abortion diagnosis of bacterial vaginosis, N. gonorrhoeae or C. trachomatis	Any antibiotic regimen; universal antibiotic prophylaxis	A placebo or nothing; or another antibiotic regimen; a screen-and-treat strategy and/or a combination of screen-and-treat and antibiotic prophylaxis	The proportion of women diagnosed with post-abortion upper genital tract infection	Other antibiotic treatments provided in the 6 weeks following the abortion Hospitalisation due to infectious complications Adverse effects of antibiotic prophylaxis or screening Proportion of women under-		Sub-group analyses were not performed; sensitivity analyses were not undertaken No data of interest to overview authors reported

**Table 2. Characteristics of included reviews** (Continued)

									going the screen-and-treat strategy who were re-infected with C. trachomatis		
CD00357	<a href="#">Mackeen 2012</a>	Techniques and materials for skin closure in caesarean section	11 studies (n = 1554)	To compare the effects of skin closure techniques and materials on maternal outcomes and time taken to perform a caesarean	Women undergoing a caesarean	Various closure techniques and materials	Different closure techniques and materials	Wound infection	Wound complications Presence of hematoma Presence of seroma Skin separation Reclosure Readmission Length of stay Pain perception Cosmesis Patient satisfaction Length of scar Total operative time Cost		Only 8 (n = 1166) of the 11 included trials contributed data: 2 studies did not report sufficiently on prespecified outcomes on which this review was focused; and 1 study did not report out-

**Table 2. Characteristics of included reviews** (Continued)

									Maternal length of hospital stay Presence of hypertrophic scar		comes separately for women undergoing caesarean. Subgroup analysis was performed; sensitivity analysis was done too
CD00951	<a href="#">Mackeen 2014<sup>a</sup></a>	Timing of intravenous prophylactic antibiotics for preventing postpartum infectious morbidity in women undergoing cesarean delivery	10 studies (n = 5041)	To compare the effects of caesarean antibiotic prophylaxis administered preoperatively versus after neonatal cord clamp on postoperative infectious complications for both the mother and the neonate	Pregnant women who have undergone caesarean delivery and received prophylactic antibiotics	Prophylactic intravenous (IV) antibiotic administration for caesarean birth 0-30 and 30-60 minutes prior to skin incision	Prophylactic antibiotic administration for caesarean birth after neonatal umbilical cord clamping	Composite maternal postpartum infectious morbidity (including serious infectious complications, endomyometritis, wound infection, or death attributed to infection)	Maternal mortality Maternal postpartum infection Placental transfer of antibiotics Breast-feeding		Subgroup analyses were not performed; sensitivity analyses were not undertaken



**Table 2. Characteristics of included reviews** (Continued)

CD01187	<a href="#">Nabhan 2016<sup>a</sup></a>	Routes of administration of antibiotic prophylaxis for preventing infection after caesarean section	10 studies (n = 1354)	To assess the benefits and harms of different routes of prophylactic antibiotics given for preventing infectious morbidity in women undergoing caesarean section	Women undergoing elective or emergency caesarean section	Prophylactic antibiotic regimens	Different route(s) of antibiotic administration	Maternal: endometritis; wound infection Infant: infant sepsis (suspected or proven)	Maternal: postpartum febrile morbidity; urinary tract infection; serious infectious complication; adverse effects of treatment on the woman; maternal length of hospital stay; readmissions Infant: oral thrush; infant length of hospital stay; immediate adverse effects of antibiotics on the infant	Combined groups of similar routes to create a single pairwise comparison	Sub-group analysis was carried out by dosage; Sensitivity analysis was performed
CD00118	<a href="#">Nelson 2014<sup>a</sup></a>	Antimicrobial prophylaxis for colorectal	260 trials (n = 43, 451) 68 differ-	To establish the effectiveness of antim-	Patients (adults and children) undergoing	All antimicrobial prophylaxis regimens	No treatment control/placebo Regi-	SSI (abdominal wound)			

**Table 2. Characteristics of included reviews** (Continued)

		surgery	ent antibiotics	crobial prophylaxis for the prevention of surgical wound infection in people undergoing colorectal surgery	either elective or emergency colorectal surgery, in which sepsis was not suspected preoperatively	delivered orally, intravenously or by intramuscular injection used to prevent postoperative infection	men differing in duration, timing, use of aerobic/anaerobic coverage, route of administration	A published gold standard regimen			
CD00526	<a href="#">Sanabria 2010<sup>a</sup></a>	Antibiotic prophylaxis for patients undergoing elective laparoscopic cholecystectomy	11 studies (n = 1664)	To assess the beneficial and harmful effects of antibiotic prophylaxis versus placebo or no prophylaxis for people undergoing elective laparoscopic cholecystectomy	Adult patients (> 17 years) undergoing laparoscopic cholecystectomy with preoperative clinical diagnosis of cholelithiasis without acute cholecystitis or other benign, non-acute	Antibiotic prophylaxis, administered intravenously or orally, prior to elective laparoscopic surgery	Placebo or no antibiotic	All-cause mortality SSI Extra-abdominal infections Adverse events Quality of life			A sensitivity analysis using worst-best case and best-worst case analyses

**Table 2. Characteristics of included reviews** (Continued)

					inflammatory disease of the gall-bladder. Jaundiced patients were excluded						
CD00376	Sanchez-Manuel 2012 <sup>a</sup>	Antibiotic prophylaxis for hernia repair	17 studies (n = 7843)	To clarify the effectiveness of antibiotic prophylaxis in reducing postoperative wound infection rates in elective open inguinal hernia repair	Adult patients undergoing open elective inguinal or femoral hernia repair, with or without the use of prosthetic material	Administration of prophylactic antibiotics	Placebo or no treatment	Wound infection rate assessed at least at 30 days after the prophylactic antibiotic treatment was given			Sensitivity analysis and subgroup analysis were conducted
CD00748	Smaill 2014 <sup>a</sup>	Antibiotic prophylaxis versus no prophylaxis for preventing infection after caesarean section	95 studies (> 15,000 women)	To assess the effects of prophylactic antibiotics compared with no prophylactic antibiotics on infectious	Women undergoing caesarean section, both elective (planned) and non-elective/emergency	Any prophylactic antibiotic regimen administered for caesarean section	Placebo or no treatment	Maternal: febrile morbidity; wound infection; endometritis; serious infectious complication	Maternal: urinary tract infection; adverse effects of treatment on the woman; length of stay in hospital		A sensitivity analysis was undertaken on the primary outcomes by study quality, omitting the 9 quasi-

**Table 2. Characteristics of included reviews** (Continued)

				complications in women undergoing caesarean section				Infant: immediate adverse effects of antibiotics on the infant; oral thrush	Infant: length of stay in hospital; long-term adverse effects; immune system development Additional outcomes: development of bacterial resistance; cost		RCTs; subgroup analyses were carried out by antibiotic regimen, type of surgery and time of administration
CD00308	<a href="#">Tanner 2006</a>	Double gloving to reduce surgical cross-infection	31 studies (n = not reported) Unit of analysis varied, gloves were collected)	To determine if additional glove protection reduces the number of surgical site or blood-borne infections in patients or the surgical team; to determine if additional	All members of the surgical team practising in a designated surgical theatre	Single gloves Double gloves Glove liners  Coloured perforation indicator systems Cloth outer gloves Steel outer gloves Triple gloves	Another/different type	Rates of SSI in surgical patients	Rates of perforations in innermost surgical gloves Rates of blood-borne infections in post-operative patients or the surgical team		2 trials were found that addressed surgical site infections in patients. Both trials reported no infections No subgroup analysis performed; sensitivity analysis was

**Table 2. Characteristics of included reviews** (Continued)

				glove protection reduces the number of perforations to the innermost pair of surgical gloves							not done
CD00292	<a href="#">Vincent 2016</a>	Disposable surgical face masks for preventing surgical wound infection in clean surgery	3 studies (n = 2106)	To determine whether the wearing of disposable surgical face masks by the surgical team during clean surgery reduces postoperative surgical wound infection	Adults and children undergoing clean surgery	The wearing, by the surgical team (scrubbed and not scrubbed), of disposable surgical face masks	No masks	The incidence of postoperative surgical wound infection	Costs Length of hospital stay Mortality rate		No subgroup analysis performed
CD00635	<a href="#">Webster 2015</a>	Use of plastic adhesive drapes during surgery for preventing surgical site	7 studies (n = 4195)	To assess the effect of adhesive drapes used during surgery on surgical	People of any age or gender, undergoing any type of inpatient or outpa-	Plastic adhesive drapes through which an incision is made (used alone or	No plastic adhesive drapes; other drapes (e.g. woven (material) or dispos-	SSI	Mortality Length of hospital stay Costs Hospital readmissions		The only subgroup analysis that was possible, based on available

**Table 2. Characteristics of included reviews** (Continued)

		infection		site infection, cost, mortality and morbidity	patient surgery	in combination with other drapes and any antiseptic skin preparation)	able (paper) drapes)		Adverse reactions Other serious infection or infectious complication such as septicaemia or septic shock		data, was of clean compared with contaminated surgery (wound classification) Sensitivity analyses were carried out by excluding trials most susceptible to bias: those with inadequate allocation concealment and uncertain or unblinded outcome assessment It was not possible to under-

**Table 2. Characteristics of included reviews** (Continued)

											take a planned sensitivity analysis based on the type of material the drape was made from due to insufficient detail about the products
CD00888	<a href="#">Wetter- slev 2015<sup>a</sup></a>	The effects of high perioperative inspiratory oxygen fraction for adult surgical patients	28 studies (n = 9330)	To assess the benefits and harms of an FIO <sub>2</sub> ≥ 60% compared with a control FIO <sub>2</sub> ≤ 40% in the perioperative setting in terms of mortality, surgical site infection, respiratory insuffi-	Surgical patients ≥ 18 years who were undergoing elective or emergency surgery	A high FIO <sub>2</sub> of ≥ 60%	A control FIO <sub>2</sub> of ≤ 40%	All-cause mortality SSI within 30 days of follow-up after surgery	All-cause mortality within 30 days of follow-up Respiratory insufficiency Serious adverse event Duration of postoperative hospitalisations Quality of life as mea-		Sub-group and sensitivity analyses were conducted, the role of bias was examined and trial sequential analysis (TSA) was applied to examine the level of evidence

**Table 2. Characteristics of included reviews** (Continued)

				ciency, serious adverse events and length of stay during the index admission for adult surgical patients					sured by the included trials		supporting or refuting a high FIO <sub>2</sub> during surgery, anaesthesia and recovery
CD00806	Wood 2016	Cyanoacrylate microbial sealants for skin preparation prior to surgery	7 studies (n = 859)	To assess the effects of the preoperative application of microbial sealants (compared with no microbial sealant) on rates of SSI in people undergoing clean surgery	Participants undergoing any type of clean surgery in an operating theatre	Microbial sealant applied to the surgical incision site immediately before surgery	No application of microbial sealant, with or without the use of traditional preoperative preparation solutions such as povidone iodine or chlorhexidine	Rates of SSI	All-cause mortality Adverse reactions Other serious infection or infectious complication Length of hospital stay Rates of hospital re-admissions Costs Postoperative antibiotic use		No subgroup nor sensitivity analysis was not done





















<sup>a</sup>This is a multi-stage review. We only extracted data from trials delivering interventions that started in the intraoperative phase



**Table 3. Characteristics of excluded reviews**

First review author + year	Reasons for exclusion
<a href="#">Cirocchi 2014</a>	Protocol
<a href="#">McCallum 2016</a>	Protocol
<a href="#">Ousey 2016</a>	Protocol
<a href="#">Smith 2016</a>	Protocol (ongoing)
<a href="#">Verschuur 2004</a>	Only included pre- and postoperative stages

**Table 4. Assessment of results by ROBIS (risk of bias in systematic reviews)**

Review title	Phase 2 (identifying concerns with the review process)				Phase 3 (forming an overall judgement of the risk of bias)
	Study eligibility criteria	Identification and selection of studies	Data collection and study appraisal	Synthesis and findings	Risk of bias in the review
Healing by primary versus secondary intention after surgical treatment for pilonidal sinus ( <a href="#">AL-Khamis 2010</a> )					
Staples versus sutures for closing leg wounds after vein graft harvesting for coronary artery bypass surgery ( <a href="#">Biancari 2010</a> )					
Perioperative glycaemic control for diabetic patients undergoing surgery ( <a href="#">Buchleitner 2012</a> )					
Warming of intravenous and irrigation fluids for pre-					

**Table 4. Assessment of results by ROBIS (risk of bias in systematic reviews) (Continued)**

venting inadvertent perioperative hypothermia (Campbell 2015)					
Scalpel versus electrosurgery for abdominal incisions (Charoenkwan 2017)	⊕	⊕	⊕	⊕	⊕
Scalpel versus no-scalpel incision for vasectomy (Cook 2014)	⊕	⊕	?	⊕	?
Tissue adhesives for closure of surgical incisions (Dumville 2014)	⊕	⊕	⊕	⊕	⊕
Preoperative skin antiseptics for preventing surgical wound infections after clean surgery (Dumville 2015)	⊕	⊕	⊕	⊕	⊕
Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery (Grocott 2012)	⊕	⊕	⊕	⊕	⊕
Different classes of antibiotics given to women routinely for preventing infection at caesarean section (Gyte 2014)	⊕	⊕	⊕	⊕	⊕
Methods of decreasing infection to improve outcomes after liver resections (	⊕	⊕	⊕	⊕	⊕

**Table 4. Assessment of results by ROBIS (risk of bias in systematic reviews) (Continued)**

Gurusamy 2011)					
An-tibiotic prophylaxis for the prevention of methicillin resistant Staphylococcus aureus (MRSA) related complications in surgical patients (Gurusamy 2013)					
Continuous versus interrupted skin sutures for non-obstetric surgery (Gurusamy 2014a)					
Subcutaneous closure versus no subcutaneous closure after non-caesarean surgical procedures (Gurusamy 2014b)					
Vaginal preparation with antiseptic solution before cesarean section for preventing postoperative infections (Haas 2014)					
Skin preparation for preventing infection following caesarean section (Hadiati 2014)					
Prophylactic antibiotics to prevent surgical site infection after breast cancer surgery (Jones 2014)					

**Table 4. Assessment of results by ROBIS (risk of bias in systematic reviews) (Continued)**

Peri-operative glycaemic control regimens for preventing surgical site infections in adults ( <a href="#">Kao 2009</a> )					
Systemic antimicrobial prophylaxis for percutaneous endoscopic gastrostomy ( <a href="#">Lipp 2013</a> )					
Perioperative antibiotics to prevent infection after first-trimester abortion ( <a href="#">Low 2012</a> )					
Techniques and materials for skin closure in caesarean section ( <a href="#">Mackeen 2012</a> )					
Timing of intravenous prophylactic antibiotics for preventing postpartum infectious morbidity in women undergoing caesarean delivery ( <a href="#">Mackeen 2014</a> )					
Routes of administration of antibiotic prophylaxis for preventing infection after caesarean section ( <a href="#">Nabhan 2016</a> )					
Antimicrobial prophylaxis for colorectal surgery ( <a href="#">Nelson 2014</a> )					

**Table 4. Assessment of results by ROBIS (risk of bias in systematic reviews)** (Continued)

Antibiotic prophylaxis for patients undergoing elective laparoscopic cholecystectomy (Sanabria 2010)					
Antibiotic prophylaxis for hernia repair (Sanchez-Manuel 2012)					
Antibiotic prophylaxis versus no prophylaxis for preventing infection after cesarean section (Smaill 2014)					
Double gloving to reduce surgical cross-infection (Tanner 2006)					
Disposable surgical face masks for preventing surgical wound infection in clean surgery (Vincent 2016)					
Use of plastic adhesive drapes during surgery for preventing surgical site infection (Webster 2015)					
The effects of high perioperative inspiratory oxygen fraction for adult surgical patients (Wetterslev 2015)					
Cyanoacrylate microbial sealants for skin preparation prior to surgery					

Table 4. Assessment of results by ROBIS (risk of bias in systematic reviews) (Continued)

(Wood 2016)											
⊕ = low risk; ⊕ = high risk; and ? = unclear risk											
Table 5.	Illustrative comparative risks (95% CI)	Risk Ratio (RR) (95% CI) # random-effects, all other RR = fixed-effect	Number of participants (studies)	Quality/certainty of the evidence (GRADE) * assessed by overview authors § assessed by review authors	GRADE Foot-note	Reported outcome values Results in brackets are RR with 95% CIs unless otherwise indicated # random-effects, all other RR = fixed-effect	Number of participants (studies)	Quality/certainty of the evidence (GRADE) * assessed by overview authors	Quality/certainty of the evidence (GRADE) * assessed by overview authors	GRADE Foot-note	Comments † meta-analysis by overview author Odd Ratio (OR)
Intervention and comparison intervention											
	Assumed risk	Corresponding risk									
	With comparator	With intervention									
1. Theatre staff attire											
1.1. Double gloving to reduce surgical cross-infection (Tanner 2006) Double latex versus double latex with	0 per 1000	0 per 1000 (0 to 0)	Not estimable	125 (2)	Low* <sup>1</sup>	N/A	N/A	N/A		N/A	Both trials reported no SSI; both trials were underpowered for this outcome

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

liners											
1.2. Dis-posable surgical face masks for preventing surgical wound infection in clean surgery (Vincent 2016) Mask versus no mask	N/A	N/A	N/A	N/A	N/A	N/A	RR 0.10 (0.01 to 1.83); RR 1.33 (0.59 to 3.02); RR 1.16 (0.73 to 1.84)	2106 (3)	Low <sup>*1,2</sup>	<sup>1</sup> Downgraded due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention) <sup>2</sup> Downgraded due to clinical heterogeneity (direction of intervention effect varied between studies)	OR reported by review authors and converted to RR by overview authors: OR 0.07 (0.00 to 1.63); OR 1.34 (0.58 to 3.07); OR 1.17 (0.70 to 1.97) Data not pooled due to clinical heterogeneity
2. Preparation of the surgical site											
2.1. Preoperative skin antiseptics for preventing surgical wound infections after clean surgery (Dumville 2015)											

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

a) 2% iodine in 90% alcohol versus 70% alcohol	13 per 1000	12 per 1000 (1 to 194)	RR 0.94 (0.06 to 14.74)	157 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	N/A	N/A	N/A	N/A	
b) Povidone-iodine (PI) paint versus soap scrub and application of methylated spirit	51 per 1000	59 per 1000 (18 to 187)	RR 1.15 (0.36 to 3.66)	200 (1)	Low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded once due to imprecision (small numbers of events)	N/A	N/A	N/A	N/A	



Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						bers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
c-1) 7.5% aqueous PI scrub/ 10% aqueous PI paint versus 10% aqueous PI paint	140 per 1000	106 per 1000 (48 to 236)	RR 0.76 (0.34 to 1.69)	178 (2)	Low* 1,2	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded once due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)**

						and harm for the inter- ven- tion)					
c-2) 7.5% aque- ous PI scrub/ 10% aque- ous PI paint versus iodophor in alcohol paint	41 per 1000	60 per 1000 (30 to 120)	RR 1.47 (0. 73 to 2. 94)	621 (6)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	N/A	N/A	N/A	N/A	2 stud- ies had no events in either group (n = 160)
c-3) 10% aque- ous PI paint versus iodophor in alcohol	20 per 1000	125 per 1000 (16 to 981)	RR 6. 25 (0.80 to 49.05)	106 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgr twice due to impre- cision (small num-	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

paint						bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)					
d-1) 7.5% aque- ous PI scrub/ 10% aque- ous PI paint versus 2% chlorhex- idine in 70% alcohol paint	N/A	N/A	N/A	N/A	N/A	N/A	N/A	100 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility	Not es- timable due to no re- ported SSI events  in either group

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

										of both benefit and harm for the inter- ven- tion)	
d-2) 10% aque- ous PI paint versus 2% chlorhex- idine in 70% alcohol paint	63 per 1000	66 per 1000 (35 to 125)	RR 1.06 (0. 56 to 2. 00)	556 (1)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	Not es- timable	100 (1)	N/A	N/A	No events in either group
d-3) Iodophor in alcohol (film- form- ing)	N/A	N/A	N/A	N/A	N/A	N/A	Not es- timable	100 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgr once due to risk of bias	No events in either group

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

paint versus 2% chlorhexidine in 70% alcohol paint										<sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	
d-4) 7.5% aqueous PI scrub followed by 10% aqueous PI paint versus 4% chlorhexidine in 70% alcohol scrub	21 per 1000	57 per 1000 (11 to 289)	RR 2.76 (0.55 to 13.86)	183 (1)	Low* <sub>1,2</sub>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded once due to imprecision (small numbers of events; wide confidence intervals)	Not estimable	127 (1)	N/A	N/A	No events in either group

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

						vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)					
d-5) 0.5% chlorhex- idine in methy- lated spirit versus PI paint	133 per 1000	62 per 1000 (36 to 109)	RR 0.47 (0. 27 to 0. 82)	542 (1)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events)	N/A	N/A	N/A	N/A	
e) 0. 75% Cholorhe idine and 1.5% cetrim- ide scrub versus 0.75% chlorhex- idine and 1.5% cetrim- ide paint	44 per 1000	44 per 1000 (6 to 296)	RR 0.98 (0. 14 to 6. 65)	91 (1)	Low* 1,2	<sup>1</sup> Down- graded once due to risk of bias <sup>2</sup> Down- graded once due to impre- cision (small num- bers of	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
f) Alcoholic solutions versus aqueous solutions	53 per 1000	41 per 1000 (27 to 62)	RR 0.77 (0.51 to 1.17)	1400 (6)	Low* 1,2	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded once due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						harm for the inter- ven- tion)					
<b>2.2. Skin preparation for preventing infection following caesarean section (Hadiati 2014)</b>											
a) . Drape versus no drape	112 per 1000	144 per 1000 (109 to 191)	RR 1.29 (0. 97 to 1. 71)	1294 (2)	Low <sup>§</sup>	§Wide confi- dence interval cross- ing the line of no ef- fect.	N/A	N/A	N/A	N/A	
b) 1- minute alcohol scrub with iodophor drape versus 5- minute iodophor scrub without drape	N/A	N/A	N/A	N/A	N/A	N/A	Not es- timable	79 (1)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit	No events in either group



**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

										and harm for the inter- ven- tion)	
c) Parachlor with iodine versus iodine alone	120 per 1000	40 per 1000 (5 to 359)	RR 0.33 (0. 04 to 2. 99)	50 (1)	Low <sup>§</sup>	<sup>§</sup> Wide confi- dence interval cross- ing the line of no ef- fect & small sample size	N/A	N/A	N/A	N/A	
d) Chlorhex idine glu- conate versus povi- done iodine	45 per 1000	95 per 1000 (9 to 974)	RR 2.10 (0.20 to 21.42)	43 (1)	Very low <sup>§</sup>	<sup>§</sup> One study with design limita- tions Wide confi- dence interval cross- ing the line of no ef- fect, few events & small sample size	N/A	N/A	N/A	N/A	
<b>2.3. Vaginal prepa- ration with anti- septic solu- tion</b>	33 per 1000	29 per 1000 (18 to 45)	RR 0.86 (0. 54 to 1. 36)	2205 (6)	Low <sup>§</sup>	<sup>§</sup> Most stud- ies con- tribut- ing data had de- sign limita-	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

before ce- sarean section for pre- venting post- oper- ative infec- tions (Haas 2014) Vaginal prepa- ration versus control						tions Wide confi- dence interval cross- ing the line of no ef- fect					
<b>2.4. Use of plastic adhesive drapes during surgery for preventing surgical site infection (Webster 2015)</b>											
a) Ad- hesive drapes versus no ad- hesive drapes	112 per 1000	138 per 1000 (114 to 166)	RR 1.23 (1. 02 to 1. 48)	3082 (5)	High <sup>§</sup>	<sup>§</sup> The total sample met require- ments for optimal infor- mation size, and the total number of events ex- ceeded 300	N/A	N/A	N/A	N/A	
b) Iodine- impreg- nated adhe- sive drapes versus no ad-	65 per 1000	67 per 1000 (43 to 104)	RR 1.03 (0. 66 to 1. 60)	1113 (2)	Moder- ate <sup>§</sup>	<sup>§</sup> There was impre- cision on at least 2 counts; the	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

hesive drapes						total sample size was too small to meet optimal infor- mation size, and the total number of events was less than 300					
2.5. Cyanoac- late micro- bial sealants for skin prepa- ration prior to surgery (Wood 2016) Micro- bial sealant versus no mi- crobial sealant	111 per 1000	59 per 1000 (27 to 130)	RR 0.53 (0. 24 to 1. 18) <sup>#</sup>	859 (7)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and	N/A	N/A	N/A	N/A	One study had no events in either group (n = 96)

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						harm for the inter- ven- tion)						
3. Making the surgical incision												
3.1. Scalpel versus electro- surgery for ma- jor ab- dom- inal in- ci- sions (	74 per 1000	79 per 1000 (55 to 114)	RR 1. 07 (0.74 to 1.54)	2178 (11)	Low <sup>§</sup>	§Serious limi- tation due to lack of infor- mation on ran- domi- sation and allo- cation con- ceal- ment in three studies con- tribut- ing more than 50% to the analysis Serious impre- cision as 95% CIs around the es- timate were wide ranging	N/A	N/A	N/A	N/A		

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						including the probability of a reduction as well as an increase in wound infection					
3.2. Scalpel versus no-scalpel incision for vasectomy (Cook 2014) No-scalpel versus standard incision	22 per 1000	7 per 1000 (2 to 21)	RR 0.31 (0.10 to 0.94)	1182 (2)	Very Low* 1,2,3	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded once due to imprecision (small numbers of events)  <sup>3</sup> Downgraded once due to heterogeneity	N/A	N/A	N/A	N/A	2 studies differed in their timing and nature of postoperative evaluations, including the evaluation of sterility; and in operator experience with the no-scalpel technique Peto OR reported

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

											by review authors and converted to RR by overview authors: Peto OR 0.34 (0.13 to 0.90)
<b>4. Treatment of patient during surgery</b>											
<b>4.1. Warming of intravenous and irrigation fluids for preventing inadvertent perioperative hypothermia (Campbell 2015)</b>	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	No SSI data provided
<b>4.2. Perioperative glycaemic control for diabetic</b>	N/A	N/A	N/A	N/A	N/A	N/A	1/10 vs. 6/22: RR 0.37 (0.05, 2.66) <sup>#</sup>	32 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgr due to imprecision (small	Only 2 included trials were categorised

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

pa- tients under- going surgery ( Buch- leitner 2012) Inten- sive ver- sus con- ven- tional gly- caemic control										num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	in intraop phase; Out- come for In- fectious compli- cations rather than SSIs RR 0.71 (0. 22 to 2. 26) <sup>†</sup>
							3/37 vs. 1/ 36; RR 2.92 (0. 32, 26. 77) <sup>#</sup>	73 (1)			<sup>2</sup> Downgr due to incon- sistency (direc- tion of inter- vention effect varied be- tween studies)
4.3. Peri-operative glycaemic control regimens for preventing surgical site infections in adults (Kao 2009)											
a) Intra- and postop- erative strict versus conven- tional gly- caemic control	N/A	N/A	N/A	N/A	N/A	N/A	1/40 vs. 2/ 38; RR 0.48 (0. 04 to 5. 03)	78 (1)	Low <sup>*1</sup>		<sup>1</sup> Downgraded twice due to impre- cision (small num- bers of events;

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

with intra-venous insulin										wide confidence interval that include the possibility of both benefit and harm for the intervention)	
b) Intraoperative strict versus conventional glycaemic control with insulin infusion	N/A	N/A	N/A	N/A	N/A	N/A	6/185 vs. 7/186; RR 0.86 (0.30 to 2.52)	371 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	Outcome for deep wound infection
c) Intra- and postoperative	N/A	N/A	N/A	N/A	N/A	N/A	0/72 vs. 9/68; RR 0.05 (0.	140 (1)	Low* <sup>1,2</sup>	<sup>1</sup> Downgraded once due to	Outcome for pneu-



**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

strict glycaemic control with intravenous glucose insulin-potassium infusion (GIK) versus conventional glycaemic control with subcutaneous insulin							00 to 0.84)			risk of bias <sup>2</sup> Downgrade once due to imprecision (small numbers of events; small sample size)	monia and wound infections
<b>4.4. Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery</b> ( <a href="#">Grocott 2012</a> ) Increased global	N/A	N/A	N/A	N/A	N/A	N/A	4/50 vs. 5/50; RR 0.80 (0.23 to 2.81)	100 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgrade once due to imprecision (small sample size)  <sup>2</sup> Downgrade once due to inconsistency	RR 0.40 (0.19 to 0.82) <sup>†</sup>

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

blood flow versus control											
							0/19 vs. 2/18; RR 0.19 (0.01 to 3.71)	37 (1)			
							3/30 vs. 8/30; RR 0.38 (0.11 to 1.28)	60 (1)			
							2/32 vs. 10/34; RR 0.21 (0.05 to 0.90)	66 (1)			
							0/30 vs. 2/60; RR 0.39 (0.02 to 7.95)	90 (1)			
4.5. The effects of high perioperative inspiratory oxygen fraction for adult surgical patients (	129 per 1000	112 per 1000 (92 to 138)	RR 0.87 (0.71 to 1.07) <sup>#</sup>	7219 (15)	Low* <sub>1,2</sub>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to heterogeneity	N/A	N/A	N/A	N/A	Review authors obtained data on (SF)-36 from the Greif 1999 trial through Daniel

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

Wetter-slev 2015)												Sessler, who was a co-author of this trial report
60% to 90% oxygen versus 30% to 40% oxygen perioperatively												
<b>5. Use of antibiotics</b>												
<b>5.1. Antibiotic prophylaxis versus no prophylaxis for preventing infection after cesarean section (intra and post)</b> (Smaill 2014) Antibiotic versus no antibiotics	89 per 1000	36 per 1000 (31 to 41)	RR 0.40 (0.35 to 0.46) <sup>#</sup>	14,407 (82)	Moderate <sup>\$</sup>	<sup>\$</sup> In most studies the assessment of bias was judged as unclear. In a third of studies the control group did not receive a placebo and lack of blinding could have influenced the assessment	N/A	N/A		N/A	N/A	2 studies had no event in either arm (n = 182) 6 out of 82 were quasi-RCTs

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						of outcomes. In less than 20% of studies was there an adequate description of sequence generation						
			RR 0.40 (0.35 to 0.46) <sup>#</sup>	12,669 (76)	N/A	N/A	N/A	N/A		N/A	N/A	Sensitivity analysis excluded the 6 quasi-RCTs; no further details available
<b>5.2. Review: Different classes of antibiotics given to women routinely for preventing infection at caesarean section (Gyte 2014)</b> Cephalosporins versus penicillins - all women												
a) Single cephalosporin versus single penicillin	33 per 1000	27 per 1000 (12 to 59)	RR 0.83 (0.38 to 1.81) <sup>#</sup>	1497 (9)	Low <sup>§</sup>	<sup>§</sup> Most studies contributing data had design limitations. Wide confidence interval cross-	N/A	N/A		N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)**

						ing the line of no ef- fect & small sample size					
b) Single cephalosporins versus peni- cillin drug combi- nation	33 per 1000	23 per 1000 (13 to 42)	RR 0.72 (0. 40 to 1. 30) <sup>#</sup>	1608 (7)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	N/A	N/A	N/A	N/A	
c) Cephalosporins drug combi- nation versus single peni-	32 per 1000	65 per 1000 (14 to 311)	RR 2.02 (0. 42 to 9. 63) <sup>#</sup>	139 (1)	Very low*1,2	<sup>1</sup> Downgr once due to risk of bias	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

cillin						<sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
d) Cephalosporin drug combination versus penicillin drug combination	37 per 1000	46 per 1000 (16 to 133)	RR 1.23 (0.42 to 3.58) <sup>#</sup>	315 (2)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals)	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						that include the possibility of both benefit and harm for the intervention)					
<b>5.3. Antibiotic prophylaxis for the prevention of methicillin-resistant Staphylococcus aureus (MRSA) related complications in surgical patients (Gurusamy 2013)</b>											
a) Pe-floxacin versus cefazolin and oxacillin (tibial fracture requiring external fixation)	90 per 1000	67 per 1000 (39 to 115)	RR 0.74 (0.43 to 1.28)	616 (1)	Very low <sup>§</sup>	<sup>§</sup> The risk of bias in the trial was high. The confidence intervals overlapped 1 and/or 0.75 and 1.25. There were fewer than 300 events in total in the intervention and control groups	N/A	N/A	N/A	N/A	Overall SSIs; Group 1: intra Group 2: intra and post
b) Er-tapenem	15 per 1000	9 per 1000	RR 0.59 (0.	672 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgr	N/A	N/A	N/A	N/A	MRSA

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

versus cefote- tan		(2 37)	to 14 to 2. 46)			once due to risk of bias					SSIs; over 30 min within 60 min prior to the ini- tial in- cision
						<sup>2</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)					
c) Cefaman- dole versus cefaman- dole and gen- tamycin	0 per 1000	0 per 1000 (0 to 0)	RR 5. 08 (0.24 to 105.24)	522 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr twice due to impre- cision (small num- bers of events;	N/A	N/A	N/A	N/A	Overall SSIs One study with 4 arms



Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
d) Cefazolin and gentamycin versus cefamandole and gentamycin	0 per 1000	0 per 1000 (0 to 0)	RR 17.67 (1.03 to 304.54)	516 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)**

						harm for the inter- ven- tion)					
e) Ce- fazolin versus cefaman- dole	8 per 1000	27 per 1000 (6 to 131)	RR 3.55 (0.75 to 16.95)	514 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	N/A	N/A	N/A	N/A	
f) Ce- fazolin versus cefa- zolin and gen- tamycin	32 per 1000	28 per 1000 (10 to 75)	RR 0.87 (0.32 to 2.36)	508 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr twice	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

						due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
g) Co-amoxiclav or cefotaxime versus placebo	375 per 1000	98 per 1000 (41 to 244)	RR 0.26 (0.11 to 0.65)	99 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgraded twice due to imprecision (small numbers of events; small sample size)	N/A	N/A	N/A	N/A	Overall SSIs
h) Vancomycin and ceftazolin versus ceftazolin (open fractures)	87 per 1000	87 per 1000 (23 to 327)	RR 1.00 (0.27 to 3.76)	92 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to	N/A	N/A	N/A	N/A	Overall SSIs

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
i) Dapto-mycin and ce-fazolin versus ce-fazolin	129 per 1000	39 per 1000 (9 to 177)	RR 0.30 (0.07 to 1.37)	113 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include	N/A	N/A	N/A	N/A	Overall SSIs; one study with 3 arms

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)**

						the possibility of both benefit and harm for the intervention)					
j) Vancomycin and cefazolin versus cefazolin (vascular surgery)	129 per 1000	125 per 1000 (49 to 323)	RR 0.97 (0.38 to 2.50)	118 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	N/A	N/A	N/A	N/A	
k) Vancomycin and cefazolin	39 per 1000	125 per 1000 (27 to 575)	RR 3.19 (0.69 to 14.65)	107 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgraded once	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

versus dapto- mycin and ce- fazolin						due to risk of bias					
						<sup>2</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)					
l) Van- comycin versus ce- furox- ime	32 per 1000	34 per 1000 (17 to 70)	RR 1.08 (0. 53 to 2. 21)	884 (1)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias	N/A	N/A	N/A	N/A	
						<sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide					

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						confidence intervals that include the possibility of both benefit and harm for the intervention)					
<b>5.4. Prophylactic antibiotics to prevent surgical site infection after breast cancer surgery (Jones 2014)</b>											
a) Pre-operative antibiotic versus placebo	N/A	N/A	N/A	N/A	N/A	N/A	3/69 vs 10/72; RR 0.31 (0.09 to 1.09)	1708 (6)	High*		RR 0.74 (0.56 to 0.98) <sup>†</sup>
							17/110 vs 19/108; RR 0.88 (0.48 to 1.60)				
							29/164 vs 32/169; RR 0.93 (0.59 to 1.47)				
							8/144 vs 13/148; RR 0.63 (0.27 to 1.48)				

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

							17/ 303 vs 26/303; RR 0.65 (0. 36 to 1. 18)					
							3/ 59 vs 5/ 59; RR 0.60 (0. 15 to 2. 40)					
b) Pre- opera- tive an- tibiotic versus none	N/A	N/A	N/A	N/A	N/A	N/A	9/ 187 vs 25/182; RR 0.35 (0. 17 to 0. 73)	987 (2)		Low* 1,2	<sup>1</sup> Downgr once due to impre- cision (small num- bers of events)  <sup>2</sup> Downgr due to incon- sistency (direc- tion of inter- vention effect varied be- tween studies)	RR 0.48 (0. 28 to 0. 82) <sup>†</sup>
							10/ 311 vs 14/307; RR 0.71 (0. 32 to 1. 56)					
c) Peri- opera- tive an- tibi- otics versus no an- tibiotic	182 per 1000	20 per 1000 (2 to 355)	RR 0.11 (0. 01 to 1. 95)	44 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded	N/A	N/A		N/A	N/A	



Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						twice due to impre- cision (small num- bers of events; wide					
<b>5.5. Sys- temic antimi- crobial pro- phy- laxis for per- cuta- neous endo- scopic gas- tros- tomy (Lipp 2013) Sys- temic antibi- otic (IV) versus placebo/ no in- terven- tion/ skin an- tiseptic</b>	242 per 1000	94 per 1000 (73 to 123)	RR 0.39 (0. 30 to 0. 51)	1271 (12)	Low* <sup>1</sup>	<sup>1</sup> Downgr twice due to risk of bias	N/A	N/A	N/A	N/A	OR re- ported by review authors and con- verted to RR by overview au- thors: OR 0.36 (0. 26 to 0. 50)
<b>5.6. Timing of intra- venous pro- phy- lactic antibi-</b>	41 per 1000	24 per 1000 (17 to 33)	RR 0.59 (0. 44 to 0. 81) <sup>#</sup>	5041 (10)	High <sup>\$</sup>		N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

otics for preventing post-partum infectious morbidity in women undergoing cesarean delivery (Mackeen 2014)											
Prophylactic intravenous antibiotics administered before cesarean incision versus after neonatal umbilical cord clamping (maternal outcomes)											
5.7. Routes of adminis-	21 per 1000	10 per 1000 (4 to 30)	RR 0.49 (0.17 to 1.43) <sup>#</sup>	859 (7)	Very low <sup>§</sup>	<sup>§</sup> Studies with design	N/A	N/A	N/A	N/A	Intra or intra & post

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

tration of antibiotic prophylaxis for preventing infection after caesarean section ( Nabhan 2016) Intra-venous (IV) ver-sus irri-gation						limita-tions Studies include rela-tively few patients and few events and have a wide 95% CI that in-cludes both appre-ciable benefit and appre-ciable harm					
<b>5.8. Antibiotic prophylaxis for pa-tients under-going elective laparo-scopic chole-cystec-tomy ( Sanabria 2010) Antibiotic prophylaxis</b>	33 per 1000	27 per 1000 (15 to 46)	RR 0.81 (0.47 to 1.42) <sup>#</sup>	1664 (11)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgr twice due to risk of bias  <sup>2</sup> Downgr once due to imprecision (small numbers of events; wide confidence intervals that	N/A	N/A	N/A	N/A	Intra or intra & post OR re-ported by review authors and con-verted to RR by overview au-thors: OR 0.87 (0.49 to 1.54) <sup>#</sup>

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

versus placebo or no prophylaxis						include the possibility of both benefit and harm for the intervention)					
5. 9. Antibiotic prophylaxis for hernia repair (Sanchez-Manuel 2012) Antibiotic prophylaxis versus placebo	46 per 1000	31 per 1000 (25 to 38)	RR 0.67 (0.54 to 0.84) <sup>#</sup>	17 (7843)	Moderate <sup>*1</sup>	<sup>1</sup> Downgraded once due to risk of bias	N/A	N/A	N/A	N/A	Intra or intra & post OR reported by review authors and converted to RR by overview authors: OR 0.64 (0.50 to 0.82) <sup>#</sup>
<b>5.10. Antimicrobial prophylaxis for colorectal surgery (Nelson 2014)</b>											
a) Antibiotic versus no antibiotic/placebo	N/A	N/A	N/A	N/A	N/A	N/A	5/49 vs 16/50; RR 0.32 (0.13 to 0.80) <sup>#</sup>	405 (5)	Low <sup>*1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to inconsistency	RR 0.25 (0.16 to 0.41) <sup>†</sup>

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

							2/30 vs 11/27; RR 0.16 (0. 04 to 0. 67) <sup>#</sup>					
							7/108 vs 8/49; RR 0.40 (0. 15 to 1. 03) <sup>#</sup>					
							3/13 vs 11/19; RR 0.40 (0. 14 to 1. 16) <sup>#</sup>					
							2/29 vs 12/31; RR 0.18 (0. 04 to 0. 73) <sup>#</sup>					
b) Du- ration of ther- apy (short- term versus long- term dura- tion an- tibi- otic)	N/A	N/A	N/A	N/A	N/A	N/A	2/31 vs 0/27; RR 4. 38 (0.22 to 87.32) <sup>#</sup>	1484 (7)		Moder- ate <sup>*1</sup>	<sup>1</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos-	RR 1.05 (0. 78 to 1. 40) <sup>†</sup> Short- term: partic- ipants who re- ceived only a single preop- erative dose; long- term: those



Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

							RR 1.00 (0. 58 to 1. 74) <sup>#</sup>					
c) Additional aerobic coverage versus no aerobic coverage	N/A	N/A	N/A	N/A	N/A	N/A	0/13 vs 3/11; RR 0.12 (0.01 to 2.14) <sup>#</sup>	230 (4)		Low* 1,2	<sup>1</sup> Downgrade once due to risk of bias	RR 0.38 (0.16 to 0.96) <sup>†#</sup>
							3/47 vs 0/50; RR 7.44 (0.39 to 140.25) <sup>#</sup>				<sup>2</sup> Downgrade once due to imprecision (small numbers of events)	
							0/26 vs 6/23; RR 0.07 (0.00 to 1.15) <sup>#</sup>					
							3/27 vs 7/33; RR 0.52 (0.15 to 1.83) <sup>#</sup>					
d) Additional anaerobic coverage versus little anaerobic coverage	N/A	N/A	N/A	N/A	N/A	N/A	19/287 vs 44/280; RR 0.42 (0.25 to 0.70) <sup>#</sup>	1098 (4)		Low* 1,2	<sup>1</sup> Downgrade once due to risk of bias	RR 0.65 (0.47 to 0.90) <sup>†#</sup>
							18/121 vs				<sup>2</sup> Downgrade once due to inconsistency	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

							16/116; RR 1.08 (0. 58 to 2. 01) <sup>#</sup>					
							7/89 vs 9/85; RR 0.74 (0. 29 to 1. 91) <sup>#</sup>					
							9/36 vs 17/84; RR 1.24 (0. 61 to 2. 51) <sup>#</sup>					
e) Oral versus intra- venous	N/A	N/A	N/A	N/A	N/A	N/A	2/35 vs 1/37; RR 2. 11 (0.20 to 22.29) <sup>#</sup>	72 (1)		Very low <sup>*1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility	



Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

										of both benefit and harm for the inter- ven- tion)	
f) Com- bined oral and intra- venous versus oral or intra- venous alone	N/A	N/A	N/A	N/A	N/A	N/A	9/ 169 vs 15/141; RR 0.50 (0. 23 to 1. 11) <sup>#</sup>	310 (1)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	
<b>5.11. Meth- ods of de- creas- ing in- fection</b>	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	No SSI data pro- vided

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

to improve outcomes after liver resections (Gurusamy 2011)												
5.12. Perioperative antibiotics to prevent infection after first-trimester abortion (Low 2012)	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A		N/A	N/A	No SSI data provided
6. Management of theatre traffic (no reviews)												
7. Wound irrigation (no reviews)												
8. Wound closure												
8.1. Continuous versus interrupted skin sutures for non-obstetric surgery (	71 per 1000	52 per 1000 (29 to 95)	RR 0.73 (0.40 to 1.33)	602 (4)	Very low <sup>§</sup>	<sup>§</sup> High risk of bias; the confidence intervals overlapped 1 and either 0.75 or 1.25, or	N/A	N/A		N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

Gu-rusamy 2014a) Con- tinu- ous ver- sus in- ter- rupted skin su- tures						both. The number of events in the inter- vention and control group was fewer than 300					
<b>8.2. Sub- cuta- neous closure versus no sub- cuta- neous closure after non- cae- sarean sur- gical proce- dures (</b> Gu-rusamy 2014b) Subcu- taneous versus no sub- cuta- neous closure	84 per 1000	71 per 1000 (44 to 112)	RR 0.84 (0. 53 to 1. 33)	815 (6)	Very low <sup>\$</sup>	<sup>\$</sup> The trial (s) was (were) of high risk of bias The confi- dence inter- vals over- lapped 1 and either 0.75 or 1.25 or both The number of events in the inter- vention and control group was fewer than 300	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

8.3. Review: Techniques and materials for skin closure in caesarean section (Mackeen 2012)										
a) Staples versus ab- sorbable subcu- ticular suture	31 per 1000	26 per 1000 (13 to 52)	RR 0.85 (0. 43 to 1. 71)	916 (6)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to incon- sistency (direc- tion of inter- vention effect varied be- tween studies)	N/A	N/A	N/A	N/A
	13 per 1000	10 per 1000 (2 to 40)	RR 0.72 (0. 17 to 3. 01)	400 (4)	Low* 1,2	<sup>1</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm	N/A	N/A	N/A	N/A  Sensi- tivity anal- ysis: Staples (1/177) versus ab- sorbable subcu- ticular suture (3/223)

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						for the inter- ven- tion) 2  Down- graded once due to incon- sistency (direc- tion of inter- ven- tion ef- fect var- ied be- tween studies)					
b) Barbed suture versus PDS suture	33 per 1000	31 per 1000 (6 to 167)	RR 0.96 (0. 18 to 5. 10)	188 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

8.4. Healing by primary versus secondary intention after surgical treatment for pilonidal sinus (AL-Khamis 2010)										
primary versus secondary intention										
a) Open versus closed (all)	77 per 1000	100 per 1000 (71 to 142)	RR 1.31 (0. 93 to 1. 85)	1231 (10)	Very low* 1,2,3	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)  <sup>3</sup> Downgr once due to incon- sistency (direc- tion of inter-	N/A	N/A	N/A	N/A

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)**

						vention effect varied be- tween studies)					
b) Closed (mid-line) versus closed (other)	33 per 1000	124 per 1000 (62 to 246)	RR 3.72 (1.86 to 7.42)	541 (5)	Low* 1,2	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr once due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	N/A	N/A	N/A	N/A	1 study out of 5 had no event in both arms
c) Classic Limberg versus modified Limberg	30 per 1000	228 per 1000 (30 to 1000)	RR 7.54 (1.00 to 57.07)	68 (1)	Very low* <sup>1,2</sup>	<sup>1</sup> Downgr once due to risk of bias  <sup>2</sup> Downgr	N/A	N/A	N/A	N/A	

Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)					
d) Karydakis versus classic Limberg	80 per 1000	260 per 1000 (91 to 743)	RR 3.25 (1.14 to 9.29)	100 (1)	Very low <sup>1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals)	N/A	N/A	N/A	N/A	



Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)

						that include the possibility of both benefit and harm for the intervention)					
<b>8.5. Staples versus sutures for closing leg wounds after vein graft harvesting for coronary artery bypass surgery (Biancari 2010)</b> Staples versus sutures	80 per 1000	97 per 1000 (48 to 192)	RR 1.20 (0.60 to 2.39)	322 (3)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	No statistically significant difference: P = 0.99	258 participants with 516 leg segments	N/A	N/A	One study was excluded from the pooled analysis due to the risk of a unit of analysis error
<b>8.6. Tissue adhesives for closure of surgical incisions (Dumville 2014)</b>											

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

a) Tissue adhesives versus sutures	23 per 1000	40 per 1000 (22 to 73)	RR 1.72 (0.94 to 3.16)	1239 (18)	Very low <sup>§</sup>	<sup>§</sup> Study 95% CI is wide; Possible unit of analysis issues	N/A	N/A	N/A	N/A	Eight studies had no events in either group (n = 495) Sensitivity analysis was conducted due to the unit of analysis issues in 3 studies; showing similar results
b) Tissue adhesives versus adhesive tape	43 per 1000	60 per 1000 (17 to 209)	RR 1.37 (0.39 to 4.81) <sup>#</sup>	190 (3)	Low <sup>§</sup>	<sup>§</sup> Study 95% CIs are very wide Evidence of inconsistency in point estimates. With the point estimate from one	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs) (Continued)**

						study lying outside the 95% CIs of another					
c) Tis- sue ad- hesives versus staples	71 per 1000	99 per 1000 (21 to 463)	RR 1.39 (0. 30 to 6. 54) <sup>#</sup>	320 (4)	Very low <sup>§</sup>	<sup>§</sup> Study 95% CIs are very wide Evi- dence of point esti- mates lying in oppo- site di- rections with the es- timate for one study lying outside the 95% CI of another	N/A	N/A	N/A	N/A	One study had no events in either group (n = 70)
d) Tis- sue ad- hesives versus other tech- niques	66 per 1000	27 per 1000 (7 to 105)	RR 0.41 (0. 11 to 1. 60) <sup>#</sup>	249 (2)	Low <sup>§</sup>	<sup>§</sup> Study 95% CIs are very wide Single study with low event rate	N/A	N/A	N/A	N/A	One study had no events in either group (n = 40)
e) Ad- hesives ver- sus ad-	47 per 1000	38 per 1000 (7 to 200)	RR 0.82 (0. 16 to 4. 31)	148 (1)	Very low <sup>§</sup>	<sup>§</sup> Study 95% CIs	N/A	N/A	N/A	N/A	

**Table 5. 'Summary of findings' table - Outcome: Surgical Site Infections (SSIs)** (Continued)

hesives: high viscos- ity ver- sus low vis- cosity						are very wide Single study with low event rate					
f) Ad- hesives versus adhe- sives: octyl- cyanoacry late versus butyl- cyanoacry late	333 per 1000	210 per 1000 (70 to 627)	RR 0.63 (0. 21 to 1. 88)	80 (2)	Low <sup>§</sup>	<sup>§</sup> The 95% CI esti- mate around the RR of 1.46 is very wide	N/A	N/A	N/A	N/A	One study had no events in either group (n = 43)
<b>9. Theatre cleansing (no reviews)</b>											
	<b>Illustrative com- parative risks (95% CI)</b> CI = confidence interval	<b>Risk Ratio (RR)</b> (95% CI)  # random- effects, all other RR =	<b>Num- ber of partici- pants (stud- ies)</b>	<b>Qual- ity/cer- tainty of the evi- dence (GRADE * as- sessed by overview</b>	<b>GRADE Foot- note</b>	<b>Re- ported out- come values Re- sults in brack- ets are RR</b>	<b>Num- ber of partici- pants (stud- ies)</b>	<b>Qual- ity/cer- tainty of the evi- dence (GRADE *as- sessed by</b>	<b>GRADE Foot- note</b>		
										Ratio (OR)	

Table 6. 'Summary of findings' table - Outcome: Mortality (Continued)

			fixed-effect		authors § assessed by review authors		with 95% CIs un- less other- wise in- dicated # ran- dom- effects, all other RR		overview authors			
	As- sumed risk	Corre- spond- ing risk										
	With com- parator	With inter- vention										
4.2. Perioperative glycaemic control for di- abetic pa- tients under- going surgery ( <a href="#">Buch- leitner 2012</a> ) Inten- sive ver- sus con- ven- tional gly- caemic control	N/A	N/A	N/A	N/A	N/A	N/A	0/10 vs. 2/22: RR 0. 42 (0.02 to 7.99) <sup>#</sup>	32 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the	RR 1.23 (0. 18 to 8. 43) <sup>†</sup>	

Table 6. 'Summary of findings' table - Outcome: Mortality (Continued)

										inter- ven- tion)		
										<sup>2</sup> Downgr once due to incon- sistency		
							2/37 vs. 0/36: RR 4. 87 (0.24 to 98.02) <sup>#</sup>	73 (1)				
4.3. Peri-operative glycaemic control regimens for preventing surgical site infections in adults (Kao 2009)												
Intra- and post- oper- ative strict versus conven- tional gly- caemic control with intra- venous insulin	N/A	N/A	N/A	N/A	N/A	N/A	6/40 vs. 7/38; RR 0. 81 (0.30 to 2.20)	78 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgr twice due to impre- cision (small num- bers of events; wide confi- dence inter- vals that include the pos- sibility of both benefit and harm for the inter- ven- tion)		
Intra- oper- ative strict	N/A	N/A	N/A	N/A	N/A	N/A	4/ 185 vs. 0/186; RR 9.	371 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgr twice		

**Table 6. 'Summary of findings' table - Outcome: Mortality** (Continued)

versus conventional glycaemic control with insulin infusion							05 (0.49 to 166.88)				due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)	
<b>4.4. Perioperative increase in global blood flow to explicit defined goals and outcomes following surgery (Grocott 2012)</b> Increased global	66 per 1000	44 per 1000 (26 to 74)	RR 0.67 (0.40 to 1.13)	1202 (15)	Low* <sup>1</sup>	<sup>1</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit	N/A	N/A	N/A	N/A	Time intervention started - Intra-operative; 3 studies have no events in either group (n= 177)	

**Table 6. 'Summary of findings' table - Outcome: Mortality (Continued)**

blood flow versus control						and harm for the intervention)						
4.5. The effects of high perioperative inspiratory oxygen fraction for adult surgical patients ( <a href="#">Wetter-slev 2015</a> ) 60% to 90% oxygen versus 30% to 40% oxygen perioperatively	164 per 1000	137 per 1000 (89 to 212)	RR 1.07 (0.87 to 1.33) <sup>#</sup>	4918 (8)	Low* <sup>1,2</sup>	<sup>1</sup> Downgraded once due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention) <sup>2</sup> Downgraded once due to heterogeneity	N/A	N/A	N/A	N/A	2 studies have no events in either group (n= 393)	
5.1. Antibiotic prophylaxis for the prevention of methicillin-resistant Staphylococcus aureus (MRSA) related complications in surgical patients ( <a href="#">Gurusamy 2013</a> )												
Co-amoxi-clav	146 per 1000	79 per 1000 (25 to 125)	RR 0.54 (0.17 to 1.61)	99 (1)	Low* <sup>1</sup>	<sup>1</sup> Downgr twice	N/A	N/A	N/A	N/A		



**Table 6. 'Summary of findings' table - Outcome: Mortality (Continued)**

or cefotaxime versus placebo		251)	72)			due to imprecision (small numbers of events; wide confidence intervals that include the possibility of both benefit and harm for the intervention)						
Vancomycin versus cefuroxime	2 per 1000	5 per 1000 (0 to 50)	RR 2.02 (0.18 to 22.18)	884 (1)	Very low <sup>*1,2</sup>	<sup>1</sup> Downgraded once due to risk of bias  <sup>2</sup> Downgraded twice due to imprecision (small numbers of events; wide confidence intervals that	N/A	N/A	N/A	N/A		

Table 6. 'Summary of findings' table - Outcome: Mortality (Continued)

						include the possibility of both benefit and harm for the intervention)						
<b>5.11. Methods of decreasing infection to improve outcomes after liver resections (Gurusamy 2011)</b>												
Long duration antibiotics versus short duration antibiotics	Not estimable	Not estimable	Not estimable	180 (1)	Very low <sup>§</sup>	<sup>§</sup> High risk of bias The number of trials were too few to assess inconsistency The confidence intervals of risk ratio overlapped 0.75 and 1.25 Publication bias could not be assessed because of the few trials	N/A	N/A	N/A	N/A	Not estimable due to no event in either group	

**Table 6. 'Summary of findings' table - Outcome: Mortality (Continued)**

Topical povidone iodine gel versus no topical povidone iodine gel	71 per 1000	96 per 1000 (17 to 538)	RR 1.35 (0.24 to 7.53)	59 (1)	Very low <sup>§</sup>	<sup>§</sup> High risk of bias The number of trials were too few to assess inconsistency The confidence intervals of risk ratio overlapped 0.75 and 1.25 Publication bias could not be assessed because of the few trials	N/A	N/A	N/A	N/A		
---	-------------	-------------------------	------------------------	--------	-----------------------	--	-----	-----	-----	-----	--	--

**Table 7. GRADE interventions according to outcomes**

Outcome	High-certainty evidence		Moderate-certainty evidence		Low-certainty evidence		Very low-certainty evidence/no studies
	What works (Important/less important benefit/harm)	What doesn't work (no important benefit/harm)	What probably works (Important/less important benefit/harm)	What probably doesn't work (no important benefit/harm)	What may work (Important/less important benefit/harm)	No current evidence of clear difference (no important benefit/harm)	Uncertainty

**Table 7. GRADE interventions according to outcomes** (Continued)

SSI	Adhesive drape (harm)	N/A	Antibiotic prophylaxis (hernia repair)	Iodine-impregnated adhesive drapes	Intra- and post-operative strict glycaemic control with intravenous glucose insulin-potassium infusion	Aqueous solutions	2% iodine in 90% alcohol
	Prophylactic intravenous antibiotics administered before caesarean incision	N/A	Antibiotic prophylaxis (caesarean section)	Duration of therapy (antimicrobial prophylaxis for colorectal surgery)	Increased global blood flow	Double gloving	Iodophor-in-alcohol paint
	Preoperative antibiotic (breast cancer surgery)	N/A		N/A	Antibiotic prophylaxis (co-amoxiclav or cefotaxime)	Disposable surgical face masks	Chlorhexidine gluconate
	N/A	N/A	N/A	N/A	N/A	PI paint	Scalpel versus electrosurgery
	N/A	N/A	N/A	N/A	Systemic antibiotic (IV)	Aqueous PI scrub	No-scalpel
	N/A	N/A	N/A	N/A	Antibiotic prophylaxis (colorectal surgery)	Chlorhexidine and cetrimide scrub	Warming of IV and irrigation fluids
	N/A	N/A	N/A	N/A	Additional aerobic coverage (colorectal surgery)	Vaginal preparation	Intensive glycaemic control
	N/A	N/A	N/A	N/A	Additional anaerobic coverage (colorectal surgery)	Skin preparation (drape; alcohol scrub with iodophor drape; parachlorometax with iodine)	Antibiotic prophylaxis (pefloxacin; ertapenem; cefamandole; cefazolin and gentamycin; cefazolin; vancomycin and cefazolin; dap-

**Table 7. GRADE interventions according to outcomes** (Continued)

							tomyacin and cefazolin; vancomycin and cefazolin; daptomycin and cefazolin; cephalosporin drug combination)
N/A	N/A	N/A	N/A	N/A	N/A	Techniques and materials for skin closure	IV versus irrigation
N/A	N/A	N/A	N/A	N/A	N/A	Microbial sealant	Antibiotic prophylaxis (elective laparoscopic cholecystectomy)
N/A	N/A	N/A	N/A	N/A	N/A	Intra- and postoperative strict glycaemic control	Oral versus intravenous (antimicrobial prophylaxis for colorectal surgery)
N/A	N/A	N/A	N/A	N/A	N/A	High perioperative inspiratory oxygen	Methods of decreasing infection to improve outcomes after liver resections
N/A	N/A	N/A	N/A	N/A	N/A	Antibiotic prophylaxis (vancomycin; single cephalosporin)	Perioperative antibiotics to prevent infection after first-trimester abortion
N/A	N/A	N/A	N/A	N/A	N/A	Combined oral and IV versus alone (antimicrobial prophylaxis for colorectal surgery)	Continuous compared with interrupted skin sutures

**Table 7. GRADE interventions according to outcomes** (Continued)

	N/A	N/A	N/A	N/A	N/A	Intraoperative strict glycaemic control	Subcutaneous closure
	N/A	N/A	N/A	N/A	N/A	N/A	Open versus closed (primary versus secondary intention)
	N/A	N/A	N/A	N/A	N/A	N/A	Staples versus sutures
	N/A	N/A	N/A	N/A	N/A	N/A	Tissue adhesives for closure
<b>Mortality</b>	N/A	N/A	N/A	N/A	N/A	Increased global blood flow	Intensive glycaemic control
	N/A	N/A	N/A	N/A	N/A	Intra- and post-operative strict or Intraoperative strict glycaemic control	Long-duration antibiotics versus short-duration antibiotics
	N/A	N/A	N/A	N/A	N/A	High perioperative inspiratory oxygen	Topical PI gel
	N/A	N/A	N/A	N/A	N/A	Antibiotic prophylaxis (co-amoxiclav or cefotaxime)	Antibiotic prophylaxis (vancomycin and cefuroxime)

**IV:** intravenous; **NA:** not applicable; **PI:** povidone iodine; **SSI:** surgical site infection

## APPENDICES

### Appendix 1. Summary of common topical antiseptics used in preoperative skin decontamination

#### Antiseptic agents

##### Alcohol

Alcohol denatures the cell wall proteins of bacteria. Alcohol rubs are usually available in preparations of 60% to 90% strength and are effective against a wide range of gram-positive and gram-negative bacteria, *Mycobacterium tuberculosis*, and many fungi and viruses. The three main alcohols used are ethanol, isopropanol and n-propanol, and some rubs may contain a mixture of these. Alcohol-based solutions usually (but not always) contain additional active ingredients to combine the rapid bacteriocidal effect of alcohol with more persistent chemical activity.

##### Iodine and iodophors

Iodine and iodophors are iodine solutions that are effective against a wide range of gram-positive and gram-negative bacteria, the tubercle bacillus (TB), fungi and viruses. These penetrate cell walls, then oxidise and substitute the microbial contents with free iodine (Hardin 1997; Mangram 1999; Warner 1988). Iodophors contain a surfactant or stabilising agent that liberates the free iodine (Wade 1980). Iodophor has largely replaced iodine as the active ingredient in antiseptics. Iodophor comprises free iodine molecules bound to a polymer such as polyvinyl pyrrolidine (i.e. povidone), so is often termed povidone iodine (PI) (Larson 1995). Typically, 10% PI formulations contain 1% available iodine (Larson 1995; Reichman 2009). PI is soluble in both water and alcohol, and available preparations include aqueous iodophor scrub and paint, aqueous iodophor one-step preparation with polymer (3M), and alcoholic iodophor with water insoluble polymer (DuraPrep).

##### Chlorhexidine

Chlorhexidine is a biguanide. It is effective against a wide range of gram-positive and gram-negative bacteria, lipophilic viruses and yeasts. Although its immediate antimicrobial activity is slower than alcohols, it is more persistent because it binds to the outermost layer of skin.

##### Triclosan

Triclosan (2,4,4'-trichloro-2'-hydroxydiphenyl ether) has been incorporated in detergents (0.4% to 1%) and alcohols (0.2% to 0.5%) used for hygienic and surgical hand antisepsis or preoperative skin disinfection. It inhibits *Staphylococci*, coliforms, enterobacteria and a wide range of gram-negative intestinal and skin flora.

### Appendix 2. Search strategy

#1MeSH descriptor: [Surgical Wound Infection] explode all trees  
#2MeSH descriptor: [Surgical Wound Dehiscence] explode all trees  
#3(surg\* near/5 infect\*):ti,ab,kw  
#4(surg\* near/5 wound\*):ti,ab,kw  
#5(surg\* near/5 site\*):ti,ab,kw  
#6(surg\* near/5 incision\*):ti,ab,kw  
#7(surg\* near/5 dehisc\*):ti,ab,kw  
#8(wound\* near/5 dehisc\*):ti,ab,kw  
#9(wound\* near/5 infect\*):ti,ab,kw  
#10(wound near/5 disruption\*):ti,ab,kw  
#11(wound next complication\*):ti,ab,kw

#12{or #1-#11}

#13 {or #1-#11} in Cochrane Reviews (Reviews and protocols)

### Appendix 3. Assessment by ROBIS signalling questions

Re-view	Study eligi-bil-ity crite-ria	Iden-tifi-ca-tion and se-lec-tion of stud-ies	Data collection and study appraisal					Synthesis and finding					Risk of bias in the review			
			Efforts were made to minimise error in data collection	Sufficient study characteristics were collected	All relevant study results were collected	Appropriate criteria to assess risk of bias	Efforts made to minimise error in risk of bias assessment	Synthesis included all studies	All pre-defined analyses followed	Synthesis was appropriate	Heterogeneity was addressed	Robust findings e.g. assessed with funnel plot or sensitivity analyses	Addressed biases in the synthesis	Interpretations of findings addressed all of the concerns identified	Relevance of identified studies to the review's research question was appropriately considered	Reviewers avoided emphasising results on the basis of their statistical significance
AL-Khami 2010	Y	Y	Y	Y	Y	Y	Y	Y	Y	PY	PY	PN	Y	PY	Y	Y
Biancari 2010	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	PN	Y	PY	Y	Y



(Continued)

<a href="#">Buch-leit-ner 2012</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	PY	Y	Y	PY	PY	Y	Y
<a href="#">Camp-bell 2015</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	PN	Y	Y
<a href="#">Charo 2017</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Cook 2014</a>	Y	Y	Y	Y	Y	N	Y	PN	PN	PN	PN	PN	PN	PN	PY	PN
<a href="#">Dumvi 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Dumvi 2015</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Gro-cott 2012</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Gyte 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Gu-rusamy 2011</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Gu-rusamy 2013</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Gu-rusamy 2014a</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Gu-rusamy 2014b</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Haas 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	PY	Y	Y

(Continued)

<a href="#">Ha-diati 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	PN	PY	PY	Y	Y
<a href="#">Jones 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Kao 2009</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	PY	PY	PN	PY	PY	Y	Y
<a href="#">Lipp 2013</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Low 2012</a>	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Mac-keen 2012</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Mac-keen 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Nab-han 2016</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	PY	Y	Y	Y	Y
<a href="#">Nel-son 2014</a>	Y	Y	Y	Y	Y	PY	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Sanabr 2010</a>	Y	Y	Y	Y	Y	PY	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Sanche</a> <a href="#">Manue 2012</a>	Y	Y	Y	Y	Y	N	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Smaill 2014</a>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
<a href="#">Tan-ner 2006</a>	Y	Y	Y	Y	Y	N	Y	Y	Y	PY	PY	N	PY	PY	Y	Y

(Continued)

Vincent 2016	Y	Y	Y	Y	Y	Y	Y	Y	Y	PY	PY	PN	PY	PY	Y	Y
Webster 2015	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Wetterslev 2015	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Wood 2016	Y	Y	Y	Y	Y	Y	Y	Y	Y	PY	Y	N	PY	PY	Y	Y
<i>Footnotes</i> Y = yes; PY = probably yes; N = no; and PN = probably no																

## CONTRIBUTIONS OF AUTHORS

**Zhenmi Liu:** conceived, designed and coordinated the overview; extracted data; analysed and interpreted data; undertook and checked quality assessment; performed statistical analysis; produced the first draft of the overview; contributed to writing and editing the overview; performed previous work that was the foundation of the current overview; wrote to study authors / experts / companies; approved the final overview prior to submission and is a guarantor of the overview.

**Jo Dumville:** conceived, designed and coordinated the overview; extracted data; checked the quality of data extraction; analysed and interpreted data; checked the quality of the statistical analysis; produced the first draft of the overview; contributed to writing and editing the overview; made an intellectual contribution to the overview; advised on the overview; secured funding; performed previous work that was the foundation of the current overview; wrote to study author / experts / companies; and approved the final overview prior to submission.

**Gill Norman:** advised on the overview; and approved the final overview prior to submission.

**Maggie Westby:** advised on the overview; and approved the final overview prior to submission.

**Jane Blazeby:** advised on the overview; secured funding; and approved the final overview prior to submission.

**Emma McFarlane:** advised on the overview; and approved the final overview prior to submission.

**Nicky Welton:** advised on the overview; and approved the final overview prior to submission.

**Louise O'Connor:** advised on the overview; and approved the final overview prior to submission.

**Julie Cawthorne:** advised on the overview; and approved the final overview prior to submission.

**Ryan George:** advised on the overview; and approved the final overview prior to submission.

**Emma Crosbie:** advised on the overview; and approved the final overview prior to submission.

**Amber Rithalia:** advised on the overview; and approved the final overview prior to submission.

**Hung-Yuan Cheng:** checked quality assessment; and approved the final overview prior to submission.

## Contributions of the editorial base:

Nicky Cullum (Co-ordinating Editor): edited the overview; advised on methodology, interpretation and overview content; approved the final overview prior to publication.

Gill Rizzello (Managing Editor): co-ordinated the editorial process; advised on content; edited the overview.

Ursula Gonthier (Editorial Assistant): edited the Plain Language Summary and reference sections.

## DECLARATIONS OF INTEREST

Zhenmi Liu: my employment at the University of Manchester is supported by a grant from the National Institute for Health Research (NIHR) UK (NIHR Systematic Review Fellowships).

Jo Dumville: I received research funding from the National Institute for Health Research (NIHR) UK for the production of systematic reviews focusing on high priority Cochrane Reviews in the prevention and treatment of wounds.

Gill Norman: my employment at the University of Manchester was funded by the National Institute for Health Research (NIHR) UK and focuses on high priority Cochrane Reviews in the prevention and treatment of wounds.

Maggie Westby: my employment at the University of Manchester was funded by the National Institute for Health Research (NIHR) UK and focuses on high priority Cochrane Reviews in the prevention and treatment of wounds.

Jane Blazeby: I receive funding from the National Institute of Health Research to undertake a feasibility study to examine whether a full trial of different types of dressing or no dressing is possible (NIHT HTA Bluebelle study).

Emma McFarlane: none known.

Nicola Welton: I have received research grants from the NIHR and the MRC. Pfizer part-fund a junior researcher working on a methodology project using historical data in a clinical area unrelated to this project. I have received honoraria from ABPI for delivering masterclasses on evidence synthesis. I have delivered a short-course on network meta-analysis to ICON plc, the funds from which were paid to my institution.

Louise O'Connor: none known.

Julie Cawthorne: none known.

Ryan George: none known.

Emma Crosbie: I am an employee of the University of Manchester and my institution has received funds in return for my membership of the CRUK Population Health Early Diagnosis Funding Panel.

Amber Rithalia: none known.

Hung-Yuan Cheng: none known.

## SOURCES OF SUPPORT

## Internal sources

- Division of Nursing, Midwifery and Social Work, School of Health Sciences University of Manchester, UK.

## External sources

- National Institute for Health Research (NIHR), UK.

This project was supported by the National Institute for Health Research, via Cochrane Infrastructure and Cochrane Programme Grant funding (NIHR Cochrane Programme Grant 13/89/08 - High Priority Cochrane Reviews in Wound Prevention and Treatment) to Cochrane Wounds. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

- National Institute for Health Research (NIHR) Systematic Review Fellowships (NIHR-RMFI-2015-06-52 Zhenmi Liu), UK.
- National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care (NIHR CLAHRC)

Greater Manchester Centre, UK.

Jo Dumville was partly funded by the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care (NIHR CLAHRC) Greater Manchester. The funder had no role in the design of the studies, data collection and analysis, decision to publish, or preparation of the manuscript. However, the review may be considered to be affiliated to the work of the NIHR CLAHRC Greater Manchester. The views expressed herein are those of the authors and not necessarily those of the NHS, NIHR or the Department of Health.

- NIHR Biomedical Centre, Bristol, UK.

This study was supported by the NIHR Biomedical Centre at the University Hospitals Bristol NHS Foundation Trust and the University of Bristol. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

- NIHR Manchester Biomedical Research Centre, UK.

This review was co-funded by the NIHR Manchester Biomedical Research Centre. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.